

Testosterone and Growth Hormone Normalization: A Retrospective Study of Health Outcomes

Enrique Ginzburg, MD^{1*}, Alvin Lin, MD², Michael Sigler, MD¹, Denise Olsen, RN²,
Nancy Klimas, MD¹, Alan Mintz, MD²

¹University of Miami Miller School of Medicine
²Cenegenics® Medical Institute, Las Vegas, Nevada

*Corresponding author

E-mail addresses: EG: EGinzbur@med.miami.edu AL: alin@cenegenics.com

Abstract

Background: Increased visceral adiposity has been associated with increased risk of coronary artery disease, Type 2 diabetes, and other types of morbidity. [1, 2, 3] Age-related declines in testosterone and growth hormone (GH) have been associated with unfavorable changes in body composition (decline of lean mass, increased adipose tissue, and decreased bone mineral density) and general health. Clinical studies to date suggest that supplementation of these hormones in patients with documented deficiencies can ameliorate the changes in body composition with the idea that these changes will reduce the known health risks associated with such changes. The changes in body mass, however, have varied with demographic differences. **A long-term retrospective study was conducted in patients who were treated with testosterone and/or GH compared with controls to evaluate treatment regimen effect on body mass and whether there are differences among these groups.**

Methods: A database survey assessed the records of 91 men and 97 women who were treated with testosterone, GH, both, or neither. The ages of these individuals range from 25 years old to 82 years old. Patients were classified into four groups based on the hormone therapy they were receiving. Patients in each group received: dehydroepiandrosterone (DHEA) but no testosterone or GH (Control; n=31; age range 40–82); testosterone only (Tes; n=17; age range 40–79); GH only (GH; n=20; age range 42–70); and testosterone plus GH (Tes+GH; n=23, age range 36–81). All patients who received testoster-

one or GH had documented diagnoses of hormone deficiencies. Prior to being examined, all patients gave blood and urine samples. Every patient also completed an extensive questionnaire in which s/he reported her/his lifetime health history, current diet, and exercise patterns. Each patient underwent an extensive physical that included cognitive evaluation, reflex and coordination testing, bone densitometry, and nutrition and exercise counseling. The retrospective evaluation of these records recorded changes in body composition, bone density, and quality of life.

Results: After a course of treatment that lasted an average of 3 years, weight decreased in women in the Control group and the Tes+GH group but remained stable in men in all treatment groups. Testosterone alone and in combination with GH produced statistically significant increases in lean mass, reductions in fat mass, and improvements in bone mineral density in both men and women. GH alone also produced these same statistically significant changes in women. Quality of life and mood improved throughout treatment for all groups. Treatments were generally safe and well tolerated.

Conclusions: In this retrospective survey of patients who were treated for clinically documented hormone deficiencies, testosterone and GH, alone or in combination, were associated with decreased percentage of adipose tissue and increased bone density in men and women across a wide age range.

Background

Since the second century, when Galen observed that castration led to declines in sexual function and general health and recommended eating animals' testicles as a way to improve vitality, medicine has progressed to a point when isolated and purified testosterone can be used clinically. Today, however, the goal is not only to enhance libido and sexual function, especially in men, but also to optimize body composition—to offset age-related hormonal changes that may contribute to reduced bone mass (osteopenia), reduced muscle mass (sarcopenia), and increased adiposity.

Testosterone levels are positively correlated with measures of body composition such as bone mineral density (BMD) and lean muscle mass. Improved BMD is associated with a decreased risk for osteoporosis and fracture. The correlation between testosterone levels and lean muscle mass is seen across a wide range of age and health status. [4] Optimal lean mass is associated with increased strength and coordination and reduced injury from falls. Reduced body fat is associated with decreased actuarial health risk, especially for coronary artery disease and type 2 diabetes mellitus.

The most obvious beneficiaries of testosterone supplementation would be hypogonadal men. All patients in the Tes

group had been diagnosed as hypogonadal when they began testosterone treatment. Because testosterone levels generally decline as men age, with free testosterone concentrations declining by about 50% between age 25 and 75, elderly men are at higher risk of becoming hypogonadal. The over-65 male population is expected to double (to over 31 million) by 2030, with the incidence of low testosterone levels increasing from 30% in the seventh decade of life to 70% in the eighth decade. [5] Accordingly, it is important to understand whether testosterone supplementation improves measures of health.

As in men, androgens in women affect body composition, mood, libido, and general well-being. Deficiency in young women may result from ovarian or adrenal dysfunction, hypothalamic amenorrhea, ovarian failure, oophorectomy, or wasting from acquired immunodeficiency syndrome. Deficiency in testosterone may occur secondary to use of estrogen, oral contraceptives, or corticosteroids. However, this condition is difficult to recognize (declining libido may be the only symptom), and the few data on testosterone therapy in premenopausal women reveal mainly the expected adverse effects of reversible hirsutism and acne; more data are needed to identify women who would be candidates for testosterone therapy and to establish therapeutically useful regimens in this population. [6]

Like patients who have low androgen levels, individuals with GH deficiency have increased adiposity. All patients in this study who received GH had clinically documented GH deficiency. Results of studies designed to determine the effects of GH on body composition in deficient adults have been mixed. Various trials have shown improvement in BMD in men, [7,8] decreased fat mass and increased lean mass in both men and women, along with significant improvements in serum lipids but not in BMD, [9] and beneficial reductions in waist-hip ratio and serum LDL cholesterol. [10]

The objective of the retrospective survey of clinical data was to assess the effects of testosterone and GH supplementation on body composition and quality of life in men and women who had been diagnosed as deficient in androgens and/or GH across a wide age range.

Methods

This study examined the records of patients treated at the Cenegenics® Medical Institute (Las Vegas, NV) during the period 1999–2006.

Patient populations: The records of 91 men were assessed in the following treatment groups: DHEA but no testosterone or GH (Control; n=31; age range 40–82); testosterone only (Tes; n=17; age range 40–79); GH only (GH; n=20; age range 42–70); and testosterone plus GH (Tes+GH; n=23, age range 36–81).

The records of 97 women were assessed in the same defined treatment groups: Control (n=27; age range 25–60); Tes (n=26; age range 38–69); GH (n=12; age range 42–71); and Tes+GH (n=32, age range 29–75).

The average length of treatment was three years.

Table 1 outlines the hormonal regimen used in male and female patients with their consent after explanation of clinical and laboratory goals to be achieved.

The main outcome measures of the study were body mass index, bone density, and quality-of-life survey results to evaluate whether treatment with testosterone and GH were statistically different than a group of patients who were not treated with these hormones and who only received DHEA supplementation. Other therapies, given adjunctively as needed to optimize clinical and laboratory parameters, included DHEA (an adrenal precursor to both estrogens and androgens), thyroid hormone, melatonin (for antioxidant and sleep-stabilizing properties), human chorionic gonadotropin in men (to facilitate weight loss and stimulate endogenous testosterone production), and estradiol and progesterone in women.

In addition to the hormonal regimens, all patients were placed on a low-glycemic diet to improve the lipid profile and increase insulin sensitivity and an exercise program to increase lean muscle mass and decrease fat mass.

All patients had comprehensive baseline and annual physical examinations as well as laboratory assessments at four- to six-month intervals. Dual x-ray absorptiometry scans were obtained annually to assess BMD at the hip and lumbar spine; fat mass and lean muscle mass were also assessed annually. Quality-of-life outcomes were rated using a standardized scale (Beck Depression Inventory and Holmes & Dickerson linear analog self-assessment scale for quality of life) to assess mood and functionality.

Data were expressed as the mean \pm standard error. Between-group differences were assessed using the Mann-Whitney U test and the Kruskal-Wallis test. For correlations, Pearson's test was used for normally distributed data; otherwise, the Spearman rank test was used. All hypothesis tests were two-tailed, with statistical significance assessed at $P < 0.05$ with 95% confidence intervals. The statistical software used was SPSS 11.5 for Windows (SPSS Inc, Chicago, IL, USA).

Results

Serum testosterone: Among men, there were no significant changes in weight (pre- and post-treatment), and there were no significant changes between the different groups (Figure 1). In the Control group, mean total testosterone increased from 545 ng/dL pretreatment to 687 ng/dL post-treatment ($P < 0.03$), and free testosterone rose from 107 to 119 pg/mL (nonsignificant). In the Tes group, the corresponding increases in total and free testosterone were 538 to 927 ng/dL ($P < 0.002$), and 94 to 167 pg/mL ($P < 0.002$). In the GH group, the increases were 596 to 707 ng/dL (nonsignificant) and 110 to 156 pg/mL ($P < 0.006$). In the Tes+GH group, the increases were 526 to 814 ng/dL ($P < 0.002$) and 88 to 126 pg/mL ($P < 0.03$).

Among women in the Tes, GH, and Tes+GH groups, increases in serum testosterone were significant but of considerably smaller magnitude.

Body weight: Among men, none of the treatment groups showed significant change in weight, whether intra group or between groups. Mean weight across all groups remained stable within the range of 192–198 lbs (87–90 kg) over the course of the trial.

However, among women, mean weight between treatment groups was more diverse at baseline (138–155 lbs, or 63–71 kg), and treatment resulted in more notable weight decreases in the Control and Tes+GH groups (4.8% and 3.2%, respectively).

Body composition: Lean mass increased significantly with Tes (3%) and Tes+GH (6%) in men, and with Tes (2%), GH (13%), and Tes+GH (3%) in women (Figure 2). Among men, the increase with Tes+GH was significantly greater than with any other regimen.

Total body fat was significantly reduced with Tes and Tes+GH in men, and with Tes, GH, and Tes+GH in women; the percentage change was substantially greater in men (Figure 3). Trunk fat was significantly reduced in the Control ($P<0.03$) and Tes+GH ($P<0.02$) groups in men, and in the Tes ($P<0.01$), GH ($P<0.04$), and Tes+GH ($P<0.01$) groups in women; truncal adiposity increased significantly in the female Control group.

BMD, measured at the hip, increased significantly with Tes (6%) and Tes+GH (3%) in men, and with Tes (5%), GH (11%), and Tes+GH (9%), as well as with the Control regimen (3%) in women (Figure 4).

Quality-of-life outcomes: Although none of the groups showed any notable abnormalities at baseline, scores on standard measures of mood, functionality, and quality of life showed improved status over the course of treatment (Table 2).

Safety: Physical and laboratory assessments and recording of adverse effects indicated that the study treatments were generally well tolerated. Among the men, mean levels of prostate specific antigen (PSA) showed statistically nonsignificant increases with all regimens (in ng/mL: from 1.05 at baseline to 1.40 post-treatment with Tes, from 1.03 to 1.20 with GH, and from 1.04 to 1.14 with Tes+GH).

Discussion

This study showed that supplementation with testosterone or GH or both, in conjunction with beneficial lifestyle changes in diet and exercise, produced statistically significant changes in measures of body composition. Treatments were generally well tolerated.

To place these findings in perspective, a systematic review and meta-analysis of 29 randomized controlled trials of testosterone therapy in over 1,000 middle-aged and aging men (mean age 64.5) found that treatment resulted in 6.2% reduction in total body fat and a 2.7% increase in lean mass, marginal improvement in strength, significant increase (3.7%) in BMD at the lumbar spine but not at the femoral neck, and reductions in cholesterol, especially in men with lower testosterone concentrations at baseline.[11] Those findings are generally consistent with the changes noted in the present study, which included men and women spanning a wider age range. However, another systematic review of the literature questioned the benefit of testosterone supplementation in men with normal testosterone levels. [12]

Testosterone supplementation by any route of administration (intramuscular, oral, or transdermal) increases BMD by increasing formation and decreasing resorption of bone, with

theoretical but as yet unproven reduction in fracture risk. [13] In a study by Snyder *et al*, individuals with lower pre-treatment serum testosterone concentration showed greater changes in lumbar spine BMD during the first three years of treatment. [14] This finding is supported by another study in which men with borderline hypogonadism showed only limited benefit from one year of testosterone therapy (reduction in body fat mass but no significant increase in BMD or lean muscle mass); however, extending the treatment might yield more robust effects. [15]

One of the most important reasons for studying the effects of testosterone treatment is that testosterone therapy also has benefit in terms of cardiovascular health. If changes in body composition measures provide a surrogate measure for decreased cardiovascular morbidity, clinicians would have a valuable tool for determining which patients to treat and guidance for determining treatment endpoints. A case-control study showed the risk of severe atherosclerotic coronary artery disease in men varied inversely with total testosterone; risk was five-fold higher among men in lowest quartile than among men in highest quartile. [16] A large-scale survey in 1,132 men ages 30–79 revealed a significant inverse relationship between blood pressure and levels of testosterone ($P<0.001$ for both systolic and diastolic pressure), but no such correlation was seen with other hormones. [17] The reduction in body fat mass associated with testosterone therapy, along with possible stabilizing effects on blood glucose, has obvious beneficial implications in terms of lowered risk of diabetes, metabolic syndrome, and cardiovascular disease. [13] It is noteworthy that the present study showed not only reductions in total body fat with testosterone and GH, but also reductions in truncal fat, which in excess is associated with an increased risk of insulin resistance.

Another potential benefit of testosterone therapy may be decreasing the risk of Alzheimer's disease. GH is believed to have neuroprotective effects, directly or in conjunction with insulin-like growth factor I (IGF-1; a polypeptide produced mainly in the liver in response to stimulation by GH). In a study of lean elderly subjects, low testosterone availability secondary to high levels of sex hormone binding globulin was associated with a higher incidence of Alzheimer's disease. [18] Although the exclusion of heavier patients may limit the applicability of these findings, and the root problem was not absolute testosterone deficiency but excessive binding leaving inadequate free testosterone, there may be a role for testosterone supplementation in patients considered at risk. The quality-of-life measurements used in the present study may provide a useful clinical measurement of cognitive function. Recent reviews on this topic concluded supplementation may minimize cognitive loss in testosterone-deficient elderly patients at risk [19] and that supplementation may be more beneficial in elderly men than in elderly women. [20]

With reference to GH, a meta-analysis of 10 randomized trials in 458 patients with GH deficiency showed that the mean change in BMD at the lumbar spine was significant at 6 and 12 months and more strongly significant at 18 and 24 months; however, the magnitude of these changes was small and of uncertain clinical relevance. [21] The idea of combining hormonal and bisphosphonate therapy is an attractive approach to improving BMD. A long-term controlled study in 30 adults with GH deficiency showed that the combination of GH and alendronate was highly effective in patients with osteoporosis. [22] In contrast, a study in 149 men showed tes-

tosterone and alendronate were comparably effective, but the combination offered no additional benefit over monotherapy with either agent. [23]

The role of DHEA supplementation to improve BMD is unclear. In a randomized controlled one-year trial in 140 men and women (ages 60–88) with low levels of sulfated DHEA at baseline, supplementation resulted in significant improvements in BMD at the spine among the women and at the hip in both sexes [24]; however, another double-blind trial in elderly men and women showed limited benefit after 2 years of treatment.[25] In this trial, the only statistically significant changes seen in the Control groups receiving DHEA were an increase in total testosterone in men and an increase in BMD at the hip in women.

One special population that merits mention is men at increased risk of prostate cancer, as there may be concern about the safety of testosterone supplementation in these patients. Although castration has been associated with regression or retardation of advanced prostate cancer, it does not automatically follow that restoration of normal testosterone levels in hypogonadal men increases the risk of carcinogenesis. In fact, a detailed review on this subject concludes that increased risk is associated with low rather than high levels of testosterone. [26] In the present study, PSA levels increased with GH and/or testosterone, but the increases were clinically as well as statistically insignificant, as all of the post-treatment values remained well within the normal range (0–2.5 ng/mL). Obviously, men with an established history of prostate cancer would not be considered candidates for treatment.

From the present study, the finding that total and free testosterone increased across all treatment groups in men, including the Control group, indicates the basic regimen of diet and exercise with DHEA and adjunctive hormonal correction as needed was also effective, although the magnitude of the increase was greater in the Tes and Tes+GH groups. Testosterone improved to target ranges with all active treatments except combined Tes+GH; in this group, the baseline level of free testosterone was unusually low and the post-treatment value, although representing a statistically significant increase, fell just short of the lower limit of the target range.

The study left several questions unanswered. Among the women, weight decreased more in the Control group than in the Tes+GH group, which strongly suggests the value of lifestyle change and the need to encourage compliance with diet and exercise; however, it is not clear why this effect was not seen in men.

Another gender issue is the unusually large increase in lean mass seen with GH therapy in women but not in men. Nor is it clear why women, but not men, on the control regimen showed an unhealthy increase in trunk fat despite the reduction in overall fat mass.

DHEA supplementation changes androgen/estrogen ratios differently in men and women, which may explain our results. There are significant increases in estrogen levels in men, but not much increase in testosterone. For women, DHEA increases androgens but does not have much effect on estrogens. [27, 28, 29]

In our study, we did find DHEA having an effect on testosterone blood levels in men.

The main limitations of this study are its design as a retrospective database survey and the lack of stratification of outcome data by demographic variables other than gender and by concurrent treatment modalities. Moreover, although diet and exercise counseling was provided to patients, it was not feasible to determine the compliance rates. Although the current findings are intriguing, they do not distinguish results in younger versus older patients, or in patients with different levels of endogenous hormones and different measures of body composition at baseline. Nevertheless, the results of this investigation are important considering the relative paucity of long-term data (follow-up >1 year) on outcomes with similar treatment strategies. As additional longer-term retrospective data become available for hormone-deficient patients, studies that evaluate correlations between administration of hormones and specific health outcomes will provide clinicians with more precise guidance regarding which patients to treat and which clinical parameters to use as treatment endpoints. By using surrogate measures such as body composition changes, clinicians may be able to more predictably reduce cardiovascular disease and cognitive decline.

Randomized controlled prospective clinical trials are planned, with larger populations followed for even longer periods, which may further clarify the proper role of hormonal supplementation as part of a comprehensive program to preserve vitality throughout life, improve identification of suitable candidates for treatment, and establish optimal individualized regimens.

Conclusions

For patients with clinically documented low androgen levels, testosterone supplementation, alone or in combination with GH (only used in patients who had been diagnosed with adult GH deficiency), produced clinically significant changes in (1) lean body mass, (2) Beck Depression test, (3) change in total body fat, and (4) BMD at the hip in both men and women across a wide age range. These results indicate that hormonal supplementation can augment lifestyle benefits, which represent physiologic responses.

Abbreviations

BMD = bone mineral density

DHEA = dehydroepiandrosterone

GH = growth hormone

PSA = prostate specific antigen

Competing interests

The Cenegenics Education and Research Foundation funded this study. Affiliated with the organization are Alvin Lin, MD, Cenegenics® Medical Institute physician; Denise Olsen, RN, manager of Cenegenics' Quality Assurance & Research Department; Alan Mintz, MD, cofounder of Cenegenics® Medical Institute.

References

1. Lebovitz HE, Banerji MA: **Point: visceral adiposity is causally related to insulin resistance.** *Diabetes Care* 2005, **28**:2322-2325.
2. Jensen MD: **Adipose tissue as an endocrine organ: implications of its distribution on free fatty acid metabolism.** *Eur Heart J Suppl* 2006, **8**, Suppl B:B13-B19.
3. Nicklas BJ, Penninx BW, et al: **Association of visceral adipose tissue with incident of myocardial infarction in older men and women.** *Am J Epidemiol* 2004, **160**(8):741-749.
4. Bross R, Casaburi R, Storer TW, Bhasin S: **Androgen effects on body composition and muscle function: implications for the use of androgens as anabolic agents in sarcopenic states.** *Baillieres Clin Endocrinol Metab* 1998, **12**:365-378.
5. Hijazi RA, Cunningham GR: **Andropause: is androgen replacement therapy indicated for the aging male?** *Annul Rev Med* 2005, **56**:117-137.
6. Kalantaridou SN, Calis KA: **Testosterone therapy in premenopausal women.** *Semin Reprod Med* 2006, **24**:106-114.
7. Snyder PJ, Biller BM, Zagar A, Jackson I, Arafah BM, Nippoldt TB, Cook DM, Mooradian AD, Kwan A, Scism-Bacon J, Chipman JJ, Hartman ML: **Effect of growth hormone replacement on BMD in adult-onset growth hormone deficiency.** *J Bone Miner Res* 2007, **22**:762-770.
8. Bravenboer N, Holzmann PJ, ter Maaten JC, Stuurman LM, Roos JC, Lips P: **Effect of long-term growth hormone treatment on bone mass and bone metabolism in growth hormone-deficient men.** *J Bone Miner Res* 2005, **20**:1778-1784.
9. Hoffman AR, Kuntze JE, Baptista J, Baum HB, Baumann GP, Biller BM, Clark RV, Cook D, Inzucchi SE, Kleinberg D, Klibanski A, Phillips LS, Ridgway EC, Robbins RJ, Schlechte J, Sharma M, Thorner MO, Vance ML: **Growth hormone (GH) replacement therapy in adult-onset gh deficiency: effects on body composition in men and women in a double-blind, randomized, placebo-controlled trial.** *J Clin Endocrinol Metab* 2004, **89**:2048-2056.
10. Franco C, Johannsson G, Bengtsson BA, Svensson J: **Baseline characteristics and effects of growth hormone therapy over two years in younger and elderly adults with adult onset GH deficiency.** *J Clin Endocrinol Metab* 2006, **91**:4408-4414.
11. Isidori AM, Giannetta E, Greco EA, Gianfrilli D, Bonifacio V, Isidori A, Lenzi A, Fabbri A: **Effects of testosterone on body composition, bone metabolism and serum lipid profile in middle-aged men: a meta-analysis.** *Clin Endocrinol (Oxf)* 2005, **63**:280-293.
12. Krause W, Mueller U, Mazur A: **Testosterone supplementation in the aging male: which questions have been answered?** *Aging Male* 2005, **8**:31-38.
13. Köhn FM: **Testosterone and body functions.** *Aging Male* 2006, **9**:183-188.
14. Snyder PJ, Peachey H, Hannoush P, Berlin JA, Loh L, Holmes JH, Dlewati A, Staley J, Santanna J, Kapoor SC, Attie MF, Haddad JG, Strom BL: **Effect of testosterone treatment on bone mineral density in men over 65 years of age.** *J Clin Endocrinol Metab* 1999, **84**:1966-1972.
15. Merza Z, Blumsohn A, Mah PM, Meads DM, McKenna SP, Wylie K, Eastell R, Wu F, Ross RJ: **Double-blind placebo-controlled study of testosterone patch therapy on bone turnover in men with borderline hypogonadism.** *Int J Androl* 2006, **29**:381-391.
16. Chute CG, Baron JA, Plymate SR, Kiel DP, Pavia AT, Lozner EC, O'Keefe T, MacDonald GJ: **Sex hormones and coronary artery disease.** *Am J Med* 1987, **83**:853-859.
17. Khaw KT, Barrett-Connor E: **Blood pressure and endogenous testosterone in men: an inverse relationship.** *J Hypertens* 1988, **6**:329-332.
18. Paoletti AM, Congia S, Lello S, Tedde D, Orrù M, Pistis M, Pilloni M, Zedda P, Loddo A, Melis GB: **Low androgenization index in elderly women and elderly men with Alzheimer's disease.** *Neurology* 2004, **62**:301-303.
19. Beauchet O: **Testosterone and cognitive function: current clinical evidence of a relationship.** *Eur J Endocrinol* 2006, **155**:773-781.
20. Hogervorst E, Bandelow S, Moffat SD: **Increasing testosterone levels and effects on cognitive functions in elderly men and women: a review.** *Curr Drug Targets CNS Neurol Disord* 2005, **4**:531-540.
21. Davidson P, Milne R, Chase D, Cooper C: **Growth hormone replacement in adults and bone mineral density: a systematic review and meta-analysis.** *Clin Endocrinol (Oxf)* 2004, **60**:92-98.
22. Biermasz NR, Hamdy NA, Pereira AM, Romijn JA, Roelfsema F: **Long-term skeletal effects of recombinant human growth hormone (rhGH) alone and rhGH combined with alendronate in GH-deficient adults: a seven-year follow-up study.** *Clin Endocrinol (Oxf)* 2004, **60**:568-575.
23. Welch BJ, Denke MA, Kermani A, Adams-Huet B, Gazmen NM, Gruntmanis U: **Comparison of testosterone, alendronate, and a combination of both therapies in men with low bone mineral density.** *J Investig Med* 2007, **55**:168-173.
24. Jankowski CM, Gozansky WS, Schwartz RS, Dahl DJ, Kittelson JM, Scott SM, Van Pelt RE, Kohrt WM: **Effects of dehydroepiandrosterone replacement therapy on bone mineral density in older adults: a randomized, controlled trial.** *J Clin Endocrinol Metab* 2006, **91**:2986-2993.
25. Nair KS, Rizza RA, O'Brien P, Dhatariya K, Short KR, Nehra A, Vittone JL, Klee GG, Basu A, Basu R, Cobelli C, Toffolo G, Dalla Man C, Tindall DJ, Melton LJ, Smith GE, Khosla S, Jensen MD: **DHEA in elderly women and**

DHEA or testosterone in elderly men. *N Engl J Med* 2006, **355**:1647-1659.

26. Raynaud JP: **Prostate cancer risk in testosterone-treated men.** *J Steroid Biochem Mol Biol* 2006, **102** (1-5):261-266.

27. Arlt W, et al: **Biotransformation of oral dehydroepiandrosterone in elderly men: significant increase in circulating estrogens.** *J Clin Endocrinol Metab* 1999; **84**(6):2170-2176.

28. Barnhart KT, et al: **The effect of dehydroepiandrosterone supplementation to symptomatic perimenopausal women on serum endocrine profiles, lipid parameters, and health-related quality of life.** *J Clin Endocrinol Metab* 1999; **84**(11):3896-3902.

29. Arlt W, et al: **Dehydroepiandrosterone supplementation in healthy men with age-related decline of dehydroepiandrosterone secretion.** *J Clin Endocrinol Metab* 2001; **86**(10):4686-4692.

Tables

Table 1. Hormonal Regimens. In addition to the use of testosterone and GH, other hormonal treatments were offered as needed to achieve normalization of clinical and laboratory status.

Table 2. Quality-of-Life Outcomes. Mood, functionality, and quality of life showed significant improvements throughout the treatment period.

Table 1.

Hormonal Regimens. In addition to the use of testosterone and GH, other hormonal treatments were offered as needed to achieve normalization of clinical and laboratory status.

Therapy (route)	Goal	Measurement
Testosterone*, men (intramuscular) <i>For Clinically Documented Hormone Deficiency</i>	Concentration ($\pm 12.5\%$) representing 66th percentile for 40-year-old men	Total 700–900 ng/dL, free 130–200 pg/mL
Testosterone, women (transdermal or sublingual) <i>For Clinically Documented Hormone Deficiency</i>	Upper 33% of normal range for premenopausal women	Total 52–70 ng/dL
Human growth hormone (subcutaneous) <i>For Clinically Documented Hormone Deficiency</i>	Upper 40% of normal range for ages 39–54 years	Rise of $\geq 100\%$ in insulin-like growth factor 1, but not above 360 ng/mL
Dehydroepiandrosterone (DHEA) (oral)	Upper 30% of normal range for young adults	Sulfated DHEA 470–619 $\mu\text{g/dL}$ in men, 280–380 $\mu\text{g/dL}$ in women

*Human chorionic gonadotropin also used as needed to increase testosterone production.

Table 2.

Quality-of-Life Outcomes. Mood, functionality, and quality of life showed significant improvements throughout the treatment period.

		Improvement by Treatment Regimen (P value)			
		Control	Tes	GH	Tes+GH
Beck Depression Inventory	Men	P<0.05	P<0.04	P<0.008	P<0.01
	Women	P<0.01	P<0.0001	NS	P<0.05
Holmes & Dickerson Scale*	Men	NS	P<0.03	NS	P<0.08
	Women	P<0.002	P<0.005	P<0.05	P<0.02

NS=non-significant.

*Linear analog self-assessment scale for quality of life.

Figures

Figure 1. Total and free testosterone increased in all treatment groups. Most of the increases were statistically significant.

Figure 2. Testosterone, alone or in combination with GH, produced significant increases in lean mass in both men and women.

Figure 3. Testosterone, alone or in combination with GH, produced significant reductions in fat mass in both men and women.

Figure 4. Testosterone, alone or in combination with GH, produced significant improvements in BMD at the hip in both men and women.

Figure 1.

Total and free testosterone increased in all treatment groups. Most of the increases were statistically significant.




Figure 2.

Testosterone, alone or in combination with GH, produced significant increases in lean mass in both men and women.




Figure 3.

Testosterone, alone or in combination with GH, produced significant reductions in fat mass in both men and women.




Figure 4.

Testosterone, alone or in combination with GH, produced significant improvements in BMD at the hip in both men and women.

