

Vitamin D deficiency: a worldwide problem with health consequences¹⁻⁴

Michael F Holick and Tai C Chen

ABSTRACT

Vitamin D deficiency is now recognized as a pandemic. The major cause of vitamin D deficiency is the lack of appreciation that sun exposure in moderation is the major source of vitamin D for most humans. Very few foods naturally contain vitamin D, and foods that are fortified with vitamin D are often inadequate to satisfy either a child's or an adult's vitamin D requirement. Vitamin D deficiency causes rickets in children and will precipitate and exacerbate osteopenia, osteoporosis, and fractures in adults. Vitamin D deficiency has been associated with increased risk of common cancers, autoimmune diseases, hypertension, and infectious diseases. A circulating level of 25-hydroxyvitamin D of >75 nmol/L, or 30 ng/mL, is required to maximize vitamin D's beneficial effects for health. In the absence of adequate sun exposure, at least 800–1000 IU vitamin D₃/d may be needed to achieve this in children and adults. Vitamin D₂ may be equally effective for maintaining circulating concentrations of 25-hydroxyvitamin D when given in physiologic concentrations. *Am J Clin Nutr* 2008;87(suppl):1080S–6S.

HISTORICAL PERSPECTIVE

Some of the earliest phytoplankton life forms on earth that have existed unchanged in the Atlantic ocean for >750 y can make vitamin D when exposed to sunlight (1, 2). Most vertebrates, including amphibians, reptiles, birds, and lower primates, depend on sun exposure for their vitamin D requirement (2). The lack of sunlight and its association with the devastating bone-deforming disease rickets in children was first recognized by Sniadecki in 1822 (3). One hundred years would pass before it was observed that exposure to ultraviolet B radiation (UVB; 290–315 nm) from a mercury arc lamp or sunlight prevented and treated rickets (4). In the early 1930s, the US government set up an agency to provide recommendations to parents about the beneficial effect of sensible exposure to sunlight for the prevention of rickets (4–6).

The fortification of milk in the 1930s with 100 IU vitamin D₂ per 8 ounces was effective in eradicating rickets in the United States and Europe. The unfortunate outbreak of hypercalcemia in the 1950s in Great Britain was blamed on the overfortification of milk with vitamin D, even though there was little evidence for this (7). Because milk was scarce at the end of the war, many local stores that sold milk would add vitamin D to it if it was not purchased by the expiration date. This was thought to extend the shelf-life of the vitamin D–fortified milk. This rise in the incidence of hypercalcemia in infants in the 1950s resulted in Europe forbidding the fortification of dairy products with vitamin D.

Only recently have Finland and Sweden begun fortifying milk with vitamin D.

SOURCES OF VITAMIN D

The major source of vitamin D for most humans is exposure to sunlight (1, 2, 4–6). As shown in **Figure 1**, seasonal variation is found in the major circulating form of vitamin D, 25-hydroxyvitamin D [25(OH)D] (8). Few foods naturally contain vitamin D, including oily fish such as salmon, mackerel, and herring and oils from fish, including cod liver oil. We recently conducted a study and observed that wild-caught salmon had on average 500–1000 IU vitamin D in 100 g (3.5 ounces), whereas farmed salmon contained ≈ 100 –250 IU vitamin D per 100-g serving (9). The most likely reason is that vitamin D is plentiful in the food chain but is not plentiful in the pelleted diet fed to farmed salmon. In the United States, milk, some juice products, some breads, yogurts, and cheeses are fortified with vitamin D. Multivitamins that contain 400 IU vitamin D and supplements containing vitamin D only are now available in various amounts including 400, 1000, 2000, 4000, 5000 and 50 000 IU vitamin D₃. The pharmaceutical form of vitamin D in the United States is vitamin D₂ and is available as 50 000 IU vitamin D₂ in a capsule or 8000 IU vitamin D₂/mL (4, 10). In Canada, Europe, Japan, and India, vitamin D₃ is available as a pharmaceutical.

CONSEQUENCES OF VITAMIN D DEFICIENCY ON THE MUSCULOSKELETAL SYSTEM

Much debate has taken place over the definition of vitamin D deficiency. Most agree that a 25(OH)D concentration <50 nmol/L, or 20 ng/mL, is an indication of vitamin D deficiency, whereas a 25(OH)D concentration of 51–74 nmol/L, or 21–29 ng/mL, is considered to indicate insufficiency; concentrations

¹ From the Department of Medicine; Section of Endocrinology, Nutrition, and Diabetes; Vitamin D, Skin and Bone Research Laboratory; Boston University Medical Center, Boston, MA.

² Presented at the symposium "Assessment of Vitamin D in Population-Based Studies," held at Experimental Biology 2007 in Washington, DC, 1 May 2007.

³ Supported in part by NIH grant M01RR00533 and by the UV Foundation.

⁴ Reprints not available. Address correspondence to MF Holick, Boston University School of Medicine, 715 Albany Street, M-1013, Boston, MA 02118. E-mail: mfholick@bu.edu.




FIGURE 1. A: Relation between hours of sunshine and serum 25-hydroxyvitamin D [25(OH)D] concentrations. ■, hours of sunshine; ●, 25(OH)D. B: Seasonal fluctuation in serum 25(OH)D according to frequency of sun exposure. ■, regular sun exposure; ◆, occasional sun exposure; ●, avoiding direct sun exposure. Adapted from reference 8.

>30 ng/mL are considered to be sufficient (10–15; **Figure 2**) This is based on the observation that intestinal calcium absorption is maximized above 80 nmol/L, or 32 ng/mL, in postmenopausal women (16) and that parathyroid hormone (PTH) concentrations in adults continue to decline and reach their nadir at ≈75–100 nmol/L, or 30–40 ng/mL (11, 14, 15). It has been assumed that children have the same requirement as adults; however, no comparable studies have been carried out on intestinal calcium transport or PTH levels in children. Vitamin D intoxication typically does not occur until 25(OH)D concentrations are >375 nmol/L, or 150 ng/mL (10, 16, 17).

Vitamin D deficiency in children will cause growth retardation (5, 18) and classic signs and symptoms of rickets (4–6, 18). In adults, vitamin D deficiency will precipitate and exacerbate both osteopenia and osteoporosis and increase the risk of fracture (10, 11, 19, 20).

Muscle weakness has long been associated with vitamin D deficiency. A vitamin D receptor is present in skeletal muscle (21), and vitamin D deficiency has been associated with proximal muscle weakness, increase in body sway, and an increased risk of falling (22–24).

Vitamin D deficiency in adults can also cause a skeletal mineralization defect. The unmineralized osteoid provides little structural support for the periosteal covering. As a result,

patients with osteomalacia often complain of isolated or global bone discomfort along with aches and pains in their joints and muscles (25–27). These patients may be misdiagnosed with fibromyalgia, dysthymia, degenerative joint disease, arthritis, chronic fatigue syndrome, and other diseases (10, 25, 28).

CAUSES OF VITAMIN D DEFICIENCY

The major source of vitamin D for humans is exposure to sunlight (4, 8, 10). Anything that diminishes the transmission of solar UVB radiation to the earth’s surface or anything that interferes with the penetration of UVB radiation into the skin will affect the cutaneous synthesis of vitamin D₃ (2, 9; **Figure 3**) Melanin is extremely efficient in absorbing UVB radiation, and,




FIGURE 2. Schematic representation of the synthesis and metabolism of vitamin D for regulating calcium, phosphorus, and bone metabolism. During exposure to sunlight, 7-dehydrocholesterol (7-DHC) in the skin is converted to previtamin D₃ (preD₃) and then by a heat-dependent process to vitamin D₃. Vitamin D (D represents D₂ or D₃) made in the skin or ingested in the diet is converted by the vitamin D-25-hydroxylase (25-OHase) to 25-hydroxyvitamin D [25(OH)D]. 25(OH)D is converted in the kidneys by the 25-hydroxyvitamin D-1α-hydroxylase (1-OHase) to its biologically active form 1,25-dihydroxyvitamin D [1,25(OH)₂D]. 1,25(OH)₂D increases the expression of the 25-hydroxyvitamin D-24-hydroxylase (24-OHase) to catabolize 1,25(OH)₂D and 25(OH)D to the water-soluble biologically inactive calcitriol acid. 1,25(OH)₂D enhances intestinal calcium absorption in the small intestine. 1,25(OH)₂D is recognized by its receptor in osteoblasts, causing an increase in the expression of receptor activator of NFκB ligand (RANKL). CaBP, calcium binding protein; ECaC, epithelial channel calcium; FGF23, fibroblast growth factor 23; OJ, orange juice; Pi, inorganic phosphate; PTH, parathyroid hormone; UVB, ultraviolet B radiation.




FIGURE 3. A schematic representation of the major causes of vitamin D deficiency and potential health consequences. AODM, adult onset diabetes mellitus; CHD, coronary heart disease; FEV₁, forced expiratory volume in 1 s; HAART, highly active antiretroviral therapy; HBP, high blood pressure; MS, multiple sclerosis; RA, rheumatoid arthritis; TB, tuberculosis; URI, urinary tract infection.

thus, increased skin pigmentation markedly reduces vitamin D₃ synthesis (29). Similarly, a sunscreen with a sun protection of 15 absorbs 99% of the incident UVB radiation, and, thus, when topically applied properly will decrease the synthesis of vitamin D₃ in the skin by 99% (30). African Americans with very dark skin have an SPF of 15, and, thus, their ability to make vitamin D in their skin is reduced by as much as 99% (9, 29). This along with decreased milk intake are the explanations for why most African Americans who live in a temperate climate are vitamin D deficient, whereas Africans living near the equator where vitamin D₃ synthesis is more efficient because of the higher flux of UVB photons are not (31, 32).

The angle at which the sun reaches the earth has a dramatic effect on the number of UVB photons that reach the earth's surface (2, 31). This is why when the zenith angle is increased during the wintertime and in the early morning and late afternoon, little if any vitamin D₃ synthesis occurs (2, 31). The practice of purdah, whereby all skin is covered and prevented from being exposed to sunlight places those who practice it at high risk of vitamin D deficiency and explains why in the sunniest areas of the world vitamin D deficiency is very common in both children and adults (33, 34). No one is immune from vitamin D deficiency. This includes both children and adults living in the United States, Europe, Middle East, India, Australia, and Asia. These studies suggest that upwards of 30–50% of children and adults are at risk of vitamin D deficiency (33–42).

Aging is associated with decreased concentrations of 7-dehydrocholesterol, the precursor of vitamin D₃ in the skin. A 70-y-old has ≈25% of the 7-dehydrocholesterol that a young adult does and thus has a 75% reduced capacity to make vitamin D₃ in the skin (43). Because vitamin D is fat soluble, it is readily taken up by fat cells. Obesity is associated with vitamin D deficiency, and it is believed to be due to the sequestration of vitamin D by the large body fat pool (44). Medications including anti-seizure medications and glucocorticoids and fat malabsorption are also common causes of deficiency (45; Figure 3).

NONSKELETAL CONSEQUENCES OF VITAMIN D DEFICIENCY

More than 80 y ago, it was reported that living at higher latitudes in the United States correlated with an increased risk of dying of common cancers (46). In the 1980s and 1990s, several observations suggested that living at higher latitudes increased the risk of developing and dying of colon, prostate, breast, and several other cancers (47–52). Because living at higher latitudes diminishes vitamin D₃ production, it was suggested that an association may exist between vitamin D deficiency and cancer mortality. Both men and women exposed to the most sunlight throughout their lives were less likely to die of cancer (50–54). Several retrospective and prospective studies that evaluated circulating concentrations of 25(OH)D support the concept that vitamin D deficiency increases the risk of developing and dying from cancer (52, 53). It has been suggested that adults with 25(OH)D of <50 nmol/L who were then followed for up to 19 y had a 30–50% increased risk of developing colorectal, breast, prostate, and many other cancers (50, 52, 54, 55). A meta-analysis showed that increasing intake of vitamin D to 1000 IU vitamin D₃/d would be associated with a decreased risk of colorectal and breast cancer of as much as 50% (53). Men who ingested >400 IU vitamin D/d had a markedly reduced risk of developing several cancers, including those of the pancreas and esophagus and non-Hodgkin lymphoma (52). Lappe et al (56) reported that postmenopausal women who received 1100 IU vitamin D₃ and 1000 mg Ca daily for 4 y reduced their risk of developing cancer by 60%.

Living at higher latitudes is associated with an increased risk of type 1 diabetes (57), multiple sclerosis (58, 59), and hypertension (60). Children who received 2000 IU vitamin D/d during the first year of life and who were followed for 31 y were found to have a reduced risk of developing type 1 diabetes by 78% compared with children who were not supplemented with vitamin D (61). Women who received >400 IU vitamin D/d were found to have a >40% reduced risk of developing multiple sclerosis (62) and rheumatoid arthritis (63). Hypertensive patients who were exposed to a tanning bed raised their blood concentrations of 25(OH)D by >180% in 3 mo and became normotensive (64). Patients who live at higher latitudes and are at risk of vitamin D deficiency are also more prone to developing schizophrenia (65), and vitamin D deficiency has been associated with depression (66). Vitamin D deficiency in pregnancy has also been associated with an increased risk of preeclampsia (67).

African Americans are at higher risk of developing and having more severe cases of tuberculosis. It has been known for >100 y that exposure to sunlight helped in the treatment of tuberculosis (68). Liu et al (69) reported that the likely mechanism is that when a macrophage is infected with tuberculosis, it stimulates the cell to increase the production of 1,25-dihydroxyvitamin D₃ [1,25(OH)₂D₃] and increase the expression of the vitamin D receptor. In combination, they enhanced the gene expression of the bacteriocidal protein cathelicidin, which is known to kill tuberculosis and other infective agents (Figure 4).

MECHANISMS OF ACTION OF VITAMIN D

Vitamin D is metabolized in the liver to 25(OH)D and then in the kidneys to 1,25(OH)₂D (70, 71; Figure 2). It is also recognized that many other tissues in the body, including




FIGURE 4. Metabolism of 25-hydroxyvitamin D [25(OH)D] to 1,25-dihydroxyvitamin D [1,25(OH)₂D] for nonskeletal functions. When a monocyte or macrophage is stimulated through its toll-like receptor 2/1 (TLR2/1) by an infective agent such as *Mycobacterium tuberculosis* (TB) or its lipopolysaccharide (LPS), the signal up-regulates the expression of vitamin D receptor (VDR) and the 25-hydroxyvitamin D-1-hydroxylase (1-OHase). 1,25(OH)₂D increases the expression of cathelicidin (CD). When 25(OH)D concentrations are ≈ 30 ng/mL, the risk of many common cancers is reduced. It is believed that the local production of 1,25(OH)₂D regulates genes that control proliferation and apoptosis. AB, B-lymphocytes; AT, T-lymphocytes; BP, blood pressure; BS, blood sugar; 24-OHase, 25-hydroxyvitamin D-24-hydroxylase; PTH, parathyroid hormone.

macrophages, brain, colon, prostate, breast, and others, have the enzymatic machinery to locally produce 1,25(OH)₂D (72–76; Figure 4). 1,25(OH)₂D produced by the kidneys enters the circulation and travels to its major target tissues the intestine and bone, where it interacts with its vitamin D receptor to enhance intestinal calcium absorption and mobilize osteoclastic activity (70; Figure 3).

The local production of 1,25(OH)₂D in non-calcium-regulating tissues such as the colon, prostate, and breast is thought to be for the purpose of regulating up to 200 genes, which helps to control cell growth and cellular differentiation and may be responsible for decreasing the risk of the cells being transformed into a malignant state (77). 1,25(OH)₂D₃ has been shown to inhibit cancer cell growth, induce cancer cell maturation, induce apoptosis, and decrease angiogenesis (77, 78; Figure 4). 1,25(OH)₂D inhibits renin production in the kidney (79) and has an immunomodulatory activity on monocytes and activated T and B lymphocytes (80–82; Figure 4).

PREVENTION AND TREATMENT OF VITAMIN D DEFICIENCY

The Institute of Medicine recommended that all children (also endorsed by the American Academy of Pediatrics) and adults up

to the age of 50 y require 200 IU vitamin D/d and adults aged 51–70 and ≥71 y need 400 and 600 IU vitamin D/d (83). The National Osteoporosis Foundation recently recommended that all postmenopausal women take 800–1000 IU vitamin D/d (84). Cheng et al (85) reported an association of low 25(OH)D concentrations with elevated serum PTH concentrations and low cortical bone density in early pubertal and prepubertal Finnish girls. This confirmed the earlier observations of Outila et al (86), who noted elevated PTH concentrations and lower forearm bone density and vitamin D deficiency in the winter in adolescent females, and Guillemant et al (87), who observed seasonal variation in PTH concentrations in growing male adolescents. When 171 prepubertal girls were given 400 IU vitamin D₂/d from October to February and 500 mg Ca supplementation, their serum 25(OH)D concentrations did not change. When these girls received 800 IU vitamin D₂/d, their blood concentrations rose during the winter but did not reach concentrations observed during the summer (88). Thus, on the basis of these and other observations, many experts now agree that in the absence of adequate sun exposure, 800–1000 IU vitamin D/d is needed for children of all ages and adults of all ages (84, 88–91), although this is not the current recommendation of pediatric or governmental organizations. Higher doses may be required if fat malabsorption, obesity, or other causes exist that would enhance vitamin D catabolism and its destruction (10, 45; Figure 2).

As many as 4 different enzymes have been suggested to be capable of converting vitamin D to 25(OH)D (92). These enzymes most likely have different K_m values for vitamin D and have different levels of negative feedback regulation by the serum 25(OH)D concentration. Thus, circulating 25(OH)D concentrations in response to vitamin D may be influenced by the baseline 25(OH)D concentration. As can be seen in Figure 5, the baseline concentration of 25(OH)D is an important factor for how a person responds to a vitamin D dose. When serum 25(OH)D concentrations were <50 nmol/L (20 ng/mL) in nursing home patients, doses of 200, 400, and 600 IU vitamin D₂/d for 5 mo (23) raised serum 25(OH)D concentrations by ≈100% to ≈62 nmol/L (24 ng/mL). Only when the dose was increased to 800 IU/d for 5 mo did concentrations rise above 75 nmol/L, or 30 ng/mL (Figure 5). However, subjects who had starting mean 25(OH)D concentrations above 64 nmol/L (25 ng/mL) showed no significant change in their serum 25(OH)D concentrations when they took 200, 400, 600, or 800 IU/d. When the baseline 25(OH)D concentration was above 50 nmol/L (20 ng/mL), only 800 IU vitamin D₂/d for 5 mo was effective in raising the serum 25(OH)D level (Figure 5). This study evaluated vitamin D₂, which has been reported to be only 30% to 50% as effective as vitamin D₃ in maintaining serum 25(OH)D concentrations (93, 94). Our data suggest that vitamin D₂ was effective in raising blood concentrations of 25(OH)D by ≥1 ng/100 IU, as has been reported for vitamin D₃ (91, 95). These data are consistent with our recent observation that 1000 IU vitamin D₂/d was as effective as 1000 IU vitamin D₃/d in raising and maintaining serum 25(OH)D concentrations (91). Thus, physiologic doses of vitamin D₂ may be equally effective as vitamin D₃ in maintaining serum 25(OH)D concentrations.

To treat vitamin D deficiency in the United States, 50 000 IU vitamin D₂ (or vitamin D₃, which is available in Canada, Europe, Japan, and India) once a week for 8 wk often attains a 25(OH)D concentration of ≈75 nmol/L (13). To maintain vitamin D sufficiency, Holick (10) recommends that 50 000 IU vitamin D₂




FIGURE 5. Mean (\pm SE) circulating concentrations (A) and changes (B) in 25-hydroxyvitamin D [25(OH)D] in nursing home residents with initial 25(OH)D serum concentrations of either <20 ng/mL (Δ – Δ) or ≥ 20 ng/mL (\bullet – \bullet) before and after receiving 0, 200, 400, 600, or 800 IU vitamin D₂/d for 5 mo from October through March. Data are from 6–16 individuals. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

every 2 wk or its equivalent will sustain 25(OH)D concentrations above 75 nmol/L.

CONCLUSION

Throughout evolution, humans have depended on the sun for their vitamin D requirement (1, 2). Indeed, a likely reason that melanin pigmentation devolved was to permit humans who migrated north and south of the equator to make enough vitamin D in their skin to satisfy their requirement (96). The recommendation for the avoidance of all sun exposure has put the world's population at risk of vitamin D deficiency (97). This has become apparent in Australia, where a dramatic increase in skin cancer rates resulted in the promotion of never exposing the skin to direct sunlight without sun protection, ie, clothing or sunscreen. The so-called sun-safe message has resulted in a marked increase in the risk of vitamin deficiency in Australia (40).

The best method for determining a person's vitamin D status is to measure a 25(OH)D concentration. Most commercial assays are reliable enough to determine a person's vitamin D status (10). These include various radioimmunoassays (98) and what is now considered to be the gold standard: liquid chromatography–tandem mass spectroscopy (14). There has been much discussion about vitamin D₂ being only ≈ 30 –50% as effective as vitamin D₃ in maintaining serum concentrations of 25(OH)D (93, 94). This, however, did not mean that vitamin D₂ was less active than vitamin D₃ once it was metabolized to 1,25(OH)₂D₂. It only meant that vitamin D₂ may need to be given in higher doses to

raise the blood concentrations of 25(OH)D above 75 nmol/L, or 30 ng/mL. Our data (Figure 5), as well as our recent observation that vitamin D₂ was as effective as vitamin D₃ in raising the blood concentrations of 25(OH)D (91), however, calls into question whether this is really necessary.

A reevaluation needs to take place of what the adequate intakes of vitamin D should be for children and adults. The literature over the past decade suggests that the Institute of Medicine recommendations in 1997 (83) are inadequate, and some experts including us suggest that both children and adults should take ≥ 800 –1000 IU vitamin D/d from dietary and supplemental sources (4, 9, 77) when sunlight is unable to provide it. This recommendation, however, has not yet been embraced either by official government or pediatric organizations in the United States, Canada, or Europe for either children or adults.

Neither of the authors had a conflict of interest.

REFERENCES

- Holick MF. Phylogenetic and evolutionary aspects of vitamin D from phytoplankton to humans. In: Pang PKT, Schreiberman MP, eds. Vertebrate endocrinology: fundamentals and biomedical implications. Vol 3. Orlando, FL: Academic Press, Inc (Harcourt Brace Jovanovich), 1989: 7–43.
- Holick MF. Vitamin D: A millennium perspective. *J Cell Biochem* 2003;88:296–307.
- Sniadecki J. Jerdrzej Sniadecki (1768–1838) on the cure of rickets. (1840) Cited by W Mozolowski. *Nature* 1939;143:121–4.
- Holick MF. Resurrection of vitamin D deficiency and rickets. *J Clin Invest* 2006;116:2062–72.
- Hess AF. Collected writings, volume I. Springfield, IL: Charles C Thomas, 1936:669–719.
- Hess AF, Unger LJ. The cure of infantile rickets by sunlight. *JAMA* 1921;77:39.
- A British Paediatric Association Report. Infantile hypercalcaemia, nutritional rickets, and infantile scurvy in Great Britain. *Br Med J* 1964;1: 1659–61.
- Brot C, Vestergaard P, Kolthoff N, Gram J, Hermann AP, Sorensen OH. Vitamin D status and its adequacy in healthy Danish perimenopausal women: relationships to dietary intake, sun exposure and serum parathyroid hormone. *Br J Nutr* 2001;86:S97–103.
- Chen TC, Chimeh F, Lu Z, et al. Factors that influence the cutaneous synthesis and dietary sources of vitamin D. *Arch Biochem Biophys* 2007;460:213–7.
- Holick MF. Vitamin D deficiency. *N Engl J Med* 2007;357:266–81.
- Chapuy MC, Schott AM, Garnero P, Hans D, Delmas PD, Meunier J. Healthy elderly French women living at home have secondary hyperparathyroidism and high bone turnover in winter. *J Clin Endocrinol Metab* 1996;81:1129–33.
- Dawson-Hughes B, Heaney RP, Holick MF, Lips P, Meunier PJ, Vieth R. Estimates of optimal vitamin D status. *Osteoporos Int* 2005;16:713–6 (editorial).
- Malabanan A, Veronikis IE, Holick MF. Redefining vitamin D insufficiency. *Lancet* 1998;351:805–6.
- Holick MF, Siris ES, Binkley N, et al. Prevalence of vitamin D inadequacy among postmenopausal North American women receiving osteoporosis therapy. *J Clin Endocrinol Metab* 2005;90:3215–24.
- Thomas KK, Lloyd-Jones DH, Thadhani RI, et al. Hypovitaminosis D in medical inpatients. *N Engl J Med* 1998;338:777–83.
- Heaney RP, Dowell MS, Hale CA, Bendich A. Calcium absorption varies within the reference range for serum 25-hydroxyvitamin D. *J Am Coll Nutr* 2003;22:142–6.
- Vieth R. Vitamin D supplementation, 25-hydroxyvitamin D concentrations, and safety. *Am J Clin Nutr* 1999;69:842–56.
- Huldshinsky K. Heilung von Rachitis durch Kunstliche Hohensonne. (Ultraviolet irradiation of rachitic children.) *Deutsche Med Wochenschr* 1919;45:712–3 (in German).
- Bakhtiyarova S, Lesnyak O, Kyznesova N, Blankenstein MA, Lips P. Vitamin D status among patients with hip fracture and elderly control subjects in Yekaterinburg, Russia. *Osteoporos Int* 2006;17:441–6.

20. Larsen ER, Mosekilde L, Foldspang A. Vitamin D and calcium supplementation prevents osteoporotic fractures in elderly community dwelling residents: a pragmatic population-based 3-year intervention study. *J Bone Miner Res* 2004;19:370–8.
21. Simpson RU, Thomas GA, Arnold AJ. Identification of 1,25-dihydroxyvitamin D₃ receptors and activities in muscle. *J Biol Chem* 1985;260:8882–91.
22. Bischoff-Ferrari HA, Dietrich T, Orav EJ, et al. Higher 25-hydroxyvitamin D concentrations are associated with better lower-extremity function in both active and inactive persons aged ≥60 y. *Am J Clin Nutr* 2004;80:752–8.
23. Broe KE, Chen TC, Weinberg J, Bischoff-Ferrari HA, Holick MF, Kiel D. A higher dose of vitamin D reduces the risk of falls in nursing home residents: a randomized, multiple-dose study. *J Am Geriatr Soc* 2007;55:234–9.
24. Visser M, Deeg DJH, Lips P. Low vitamin D and high parathyroid hormone levels as determinants of loss of muscle strength and muscle mass (sarcopenia): the longitudinal aging study Amsterdam. *J Clin Endocrinol Metab* 2003;88:5766–72.
25. Malabanan AO, Turner AK, Holick MF. Severe generalized bone pain and osteoporosis in a premenopausal black female: effect of vitamin D replacement. *J Clin Densitom* 1998;1:201–204.
26. Aaron JE, Gallagher JC, Anderson J, et al. Frequency of osteomalacia and osteoporosis in fractures of the proximal femur. *Lancet* 1974;1:229–33.
27. Al-Ali H, Fuleihan GEH. Nutritional osteomalacia: substantial clinical improvement and gain in bone density post therapy. *J Clin Densitom* 2000;3:97–101.
28. Plotnikoff GA, Quigley JM. Prevalence of severe hypovitaminosis D in patients with persistent, nonspecific musculoskeletal pain. *Mayo Clin Proc* 2003;78:1463–70.
29. Clemens TL, Adams JS, Henderson SL, Holick MF. Increased skin pigment reduces the capacity of skin to synthesize vitamin D₃. *Lancet* 1982;74–76.
30. Matsuoka LY, Ide L, Wortsman J, MacLaughlin JA, Holick MF. Sunscreens suppress cutaneous vitamin D₃ synthesis. *J Clin Endocrinol Metab* 1987;64:1165–8.
31. Webb AR, Kline L, Holick MF. Influence of season and latitude on the cutaneous synthesis of vitamin D₃: exposure to winter sunlight in Boston and Edmonton will not promote vitamin D₃ synthesis in human skin. *J Clin Endocrinol Metab* 1988;67:373–8.
32. Thacher TD, Fischer PR, Strand MA, Pettifor JM. Nutritional rickets around the world: causes and future directions. *Ann Trop Paediatr* 2006;26:1–16.
33. Sedrani SH. Low 25-hydroxyvitamin D and normal serum calcium concentrations in Saudi Arabia: Riyadh region. *Ann Nutr Metab* 1984;28:181–5.
34. Fuleihan GEH, Nabulsi M, Choucair M, et al. Hypovitaminosis D in healthy schoolchildren. *Pediatrics* 2001;107:53–9.
35. Tangpricha V, Pearce EN, Chen TC, Holick MF. Vitamin D insufficiency among free-living healthy young adults. *Am J Med* 2002;112:659–62.
36. Gordon CM, DePeter KC, Estherann G, Emans SJ. Prevalence of vitamin D deficiency among healthy adolescents. *Arch Pediatr Adolesc Med* 2004;158:531–7.
37. Sullivan SS, Rosen CJ, Halteman WA, Chen TC, Holick MF. Adolescent girls in Maine at risk for vitamin D insufficiency. *J Am Diet Assoc* 2005;105:971–4.
38. Lips P, Duong T, Oleksik A, et al. A global study of vitamin D status and parathyroid function in postmenopausal women with osteoporosis: baseline data from the multiple outcomes of raloxifene evaluation clinical trial. *J Clin Endocrinol Metab* 2001;86:1212–21.
39. Marwaha RK, Tandon N, Reddy D, et al. Vitamin D and bone mineral density status of healthy schoolchildren in northern India. *Am J Clin Nutr* 2005;82:477–82.
40. McGrath JJ, Kimlin MG, Saha S, Eyles DW, Parisi AV. Vitamin D insufficiency in south-east Queensland. *Med J Aust* 2001;174:150–1.
41. Sato Y, Iwamoto J, Kanoko T, Satoh K. Amelioration of osteoporosis and hypovitaminosis D by sunlight exposure in hospitalized, elderly women with Alzheimer's disease: a randomized controlled trial. *J Bone Miner Res* 2005;20:1327–33.
42. Nesby-O'Dell S, Scanlon KS, Cogswell ME, Gillespie C, Hollis BW, Looker AC. Hypovitaminosis D prevalence and determinants among African American and white women of reproductive age: third National Health and Nutrition Examination Survey, 1988–1994. *Am J Clin Nutr* 2002;76:187–92.
43. Holick MF, Matsuoka LY, Wortsman J. Age, vitamin D, and solar ultraviolet. *Lancet* 1989;2(8671):1104–5.
44. Wortsman J, Matsuoka LY, Chen TC, Lu Z, Holick MF. Decreased bioavailability of vitamin D in obesity. *Am J Clin Nutr* 2000;72:690–3.
45. Zhou C, Assem M, Tay JC, et al. Steroid and xenobiotic receptor and vitamin D receptor crosstalk mediates CYP24 expression and drug-induced osteomalacia. *J Clin Invest* 2006;116:1703–12.
46. Apperly FL. The relation of solar radiation to cancer mortality in North America. *Cancer Res* 1941;1:191–5.
47. Garland CF, Comstock GW, Garland FC, Helsing KJ, Shaw EK, Gorham ED. Serum 25-hydroxyvitamin D and colon cancer: eight-year prospective study. *Lancet* 1989;2:1176–8.
48. Gorham ED, Garland CF, Garland FC, et al. Vitamin D and prevention of colorectal cancer. *J Steroid Biochem Mol Biol* 2005;97:179–94.
49. Hanchette CL, Schwartz GG. Geographic patterns of prostate cancer mortality. *Cancer* 1992;70:2861–9.
50. Grant WB. An estimate of premature cancer mortality in the U.S. due to inadequate doses of solar ultraviolet-B radiation. *Cancer* 2002;94:1867–75.
51. Grant WG, Garland CF. The association of solar ultraviolet B (UVB) with reducing risk of cancer: multifactorial ecologic analysis of geographic variation in age-adjusted cancer mortality rates. *Anticancer Res* 2006;26:2687–700.
52. Giovannucci E, Liu Y, Rimm EB, et al. Prospective study of predictors of vitamin D status and cancer incidence and mortality in men. *J Natl Cancer Inst* 2006;98:451–9.
53. Garland CF, Garland FC, Gorham ED, et al. The role of vitamin D in cancer prevention. *Am J Public Health* 2006;96:252–61.
54. Ahonen MH, Tenkanen L, Teppo L, Hakama M, Tuohimaa P. Prostate cancer risk and prediagnostic serum 25-hydroxyvitamin D levels (Finland). *Cancer Causes Control* 2000;11:847–52.
55. Garland C, Shekelle RB, Barrett-Connor E, Criqui MH, Ross AH, Oglesby P. Dietary vitamin D and calcium and risk of colorectal cancer: a 19-year prospective study in men. *Lancet* 1985;9:307–9.
56. Lappe JM, Travers-Gustafson D, Davies KM, Recker RR, Heaney RP. Vitamin D and calcium supplementation reduces cancer risk: results of a randomized trial. *Am J Clin Nutr* 2007;85:1586–91.
57. Stene LC, Ulriksen J, Magnus P, Joner G. Use of cod liver oil during pregnancy associated with lower risk of type I diabetes on the offspring. *Diabetologia* 2000;43:1093–8.
58. Ponsoy A-L, McMichael A, van der Mei I. Ultraviolet radiation and autoimmune disease: insights from epidemiological research. *Toxicology* 2002;181–182:71–8.
59. Embry AF, Snowdon LR, Vieth R. Vitamin D and seasonal fluctuations of gadolinium-enhancing magnetic resonance imaging lesions in multiple sclerosis. *Ann Neurol* 2000;48:271–2.
60. Rostand SG. Ultraviolet light may contribute to geographic and racial blood pressure differences. *Hypertension* 1979;30:150–6.
61. Hypponen E, Laara E, Jarvelin M-R, Virtanen SM. Intake of vitamin D and risk of type 1 diabetes: a birth-cohort study. *Lancet* 2001;358:1500–3.
62. Munger KL, Zhang SM, O'Reilly E, et al. Vitamin D intake and incidence of multiple sclerosis. *Neurology* 2004;62:60–5.
63. Merlino LA, Curtis J, Mikuls TR, Cerhan JR, Criswell LA, Saag KG. Vitamin D intake is inversely associated with rheumatoid arthritis. *Arthritis Rheum* 2004;50:72–7.
64. Krause R, Buhning M, Hopfenmuller W, Holick MF, Sharma AM. Ultraviolet B and blood pressure. *Lancet* 1998;352(9129):709–10.
65. McGrath J, Selden JP, Chant D. Long-term trends in sunshine duration and its association with schizophrenia birth rates and age at first registration—data from Australia and the Netherlands. *Schizophr Res* 2002;54:199–212.
66. Gloth FM III, Alam W, Hollis B. Vitamin D vs. broad spectrum phototherapy in the treatment of seasonal affective disorder. *J Nutr Health Aging* 1999;3:5–7.
67. Bodnar LM, Catov JM, Simhan HN, Holick MF, Powers RW, Roberts JM. Maternal vitamin D deficiency increases the risk of preeclampsia. *J Clin Endocrinol Metab* 2007;92:3517–22.

68. Chan TYK. Vitamin D deficiency and susceptibility to tuberculosis. *Calcif Tissue Int* 2000;66:476–8.
69. Liu PT, Stenger S, Li H, et al. Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. *Science* 2006;317:1770–3.
70. Holick MF, Garabedian M. Vitamin D: photobiology, metabolism, mechanism of action, and clinical applications. In: Favus MJ, ed. *Primer on the metabolic bone diseases and disorders of mineral metabolism*, 6th ed. Washington, DC: American Society for Bone and Mineral Research, 2006:129–37.
71. DeLuca H. Overview of general physiologic features and functions of vitamin D. *Am J Clin Nutr* 2004;80(suppl):1689S–96S.
72. Bikle DD. Vitamin D: role in skin and hair. In: Feldman D, ed. *Vitamin D*. Burlington, MA: Elsevier Academic Press, 2005:609–30.
73. Cross HS, Bareis P, Hofer H, Bischof MG, Bajna E, Kriwanek S. 25-Hydroxyvitamin D₃-1-hydroxylase and vitamin D receptor gene expression in human colonic mucosa is elevated during early cancerogenesis. *Steroids* 2001;66:287–92.
74. Tangpricha V, Flanagan JN, Whitlatch LW, et al. 25-Hydroxyvitamin D-1 α -hydroxylase in normal and malignant colon tissue. *Lancet* 2001;357:1673–4.
75. Schwartz GG, Whitlatch LW, Chen TC, Lokeshwar BL, Hollick MF. Human prostate cells synthesize 1,25-dihydroxyvitamin D₃ from 25-hydroxyvitamin D₃. *Cancer Epidemiol Biomarkers Prev* 1998;7:391–5.
76. Mawer EB, Hayes ME, Heys SE, et al. Constitutive synthesis of 1,25-dihydroxyvitamin D₃ by a human small cell lung cell line. *J Clin Endocrinol Metab* 1994;79:554–60.
77. Nagpal S, Na S, Rathnachalam R. Noncalcemic actions of vitamin D receptor ligands. *Endocr Rev* 2005;26:662–87.
78. Mantell DJ, Owens PE, Bundred NJ, Mawer EB, Canfield AE. 1 α ,25-dihydroxyvitamin D₃ inhibits angiogenesis in vitro and in vivo. *Circ Res* 2000;87:214–20.
79. Li YC. Vitamin D regulation of the renin-angiotensin system. *J Cell Biochem* 2003;88:327–31.
80. Mathieu C, Adorini L. The coming of age of 1,25-dihydroxyvitamin D₃ analogs as immunomodulatory agents. *Trends Mol Med* 2002;8:174–9.
81. Adorini L. 1,25-Dihydroxyvitamin D₃ analogs as potential therapies in transplantation. *Curr Opin Investig Drugs* 2002;3:1458–63.
82. Cantorna MT, Zhu Y, Froicu M, Wittke A. Vitamin D status, 1,25-dihydroxyvitamin D₃, and the immune system. *Am J Clin Nutr* 2004;80(suppl):1717S–20S.
83. Standing Committee on the Scientific Evaluation of Dietary Reference Intakes, Food and Nutrition Board, Institute of Medicine. *Dietary reference intakes for calcium, phosphorus, magnesium, vitamin D and fluoride*. Washington, DC: National Academy Press, 1999.
84. National Osteoporosis Foundation. Statement on calcium and vitamin D. Internet: <http://www.nof.org/prevention/calcium/htm> (accessed 16 November 2007).
85. Cheng S, Tylavsky F, Kroger H, et al. Association of low 25-hydroxyvitamin D concentrations with elevated parathyroid hormone concentrations and low cortical bone density in early pubertal and prepubertal Finnish girls. *Am J Clin Nutr* 2003;78:485–92.
86. Outila TA, Karkkainen MU, Lamberg-Allardt CJ. Vitamin D status affects serum parathyroid hormone concentrations during winter in female adolescents: associations with forearm bone mineral density. *Am J Clin Nutr* 2001;74:206–10.
87. Guillemant J, Cabrol S, Allemandou A, Peres G, Guillemant S. Vitamin D-dependent seasonal variation of PTH in growing male adolescents. *Bone* 1995;17:513–6.
88. Lehtonen-Veromaa M, Mottonen T, Nuotio I, Irjala K, Viikari J. The effect of conventional vitamin D₂ supplementation on serum 25(OH)D concentration is weak among peripubertal Finnish girls: a 3-yr prospective study. *Eur J Clin Nutr* 2002;56:431–7.
89. Tangpricha V, Koutkia P, Rieke SM, Chen TC, Perez AA, Holick MF. Fortification of orange juice with vitamin D: a novel approach to enhance vitamin D nutritional health. *Am J Clin Nutr* 2003;77:1478–83.
90. Vieth R, Garland C, Heaney R, et al. The urgent need to reconsider recommendations for vitamin D nutrition intake. *Am J Clin Nutr* 2007;85:649–50.
91. Holick MF, Biancuzzo RM, Chen TC, et al. Vitamin D₂ is as effective as vitamin D₃ in maintaining circulating concentrations of 25-hydroxyvitamin D. *J Clin Endocrinol Metab* 2007 Dec 18 [Epub ahead of print].
92. Jones G. Expanding role for vitamin D in chronic kidney disease: importance of blood 25-OH-D levels and extra-renal 1 α -hydroxylase in the classical and nonclassical actions of 1 α ,25-dihydroxyvitamin D₃. *Semin Dial* 2007;20:316–24.
93. Tang HM, Cole DEC, Rubin LA, Pierratos A, Siu S, Vieth R. Evidence that vitamin D₃ increases serum 25-hydroxyvitamin D more efficiently than does vitamin D₂. *Am J Clin Nutr* 1998;68:854–8.
94. Armas LAG, Hollis B, Heaney RP. Vitamin D₂ is much less effective than vitamin D₃ in humans. *J Clin Endocrinol Metab* 2004;89:5387–91.
95. Heaney RP, Davies KM, Chen TC, Holick MF, Barger-Lux MJ. Human serum 25-hydroxycholecalciferol response to extended oral dosing with cholecalciferol. *Am J Clin Nutr* 2003;77:204–10.
96. Loomis WF. Vitamin D, sunlight and natural selection. *Science* 1967;157:501–4.
97. Wolpowitz D, Gilchrist BA. The vitamin D questions: how much do you need and how should you get it? *J Am Acad Dermatol* 2006;54:301–17.
98. Binkley N, Krueger D, Cowgill CS et al. Assay variation confounds the diagnosis of hypovitaminosis D: a call for standardization. *J Clin Endocrinol Metab* 2004;89:3152–7.