

Review Article

Link between PI3K/AKT/PTEN Pathway and NOX Protein in Diseases

Atsuko Nakanishi[#], Yoko Wada, Yasuko Kitagishi, Satoru Matsuda^{#,*}

Department of Food Science and Nutrition, Nara Women's University, Kita-Uoya Nishimachi, Nara 630-8506, Japan

[Received September 30, 2013; Revised October 27, 2013; Accepted October 30, 2013]

ABSTRACT: Accumulating evidence has revealed that the PI3K/AKT/PTEN pathway acts as a pivotal determinant of cell fate regarding senescence and apoptosis, which is mediated by intracellular reactive oxygen species (ROS) generation. NADPH oxidase (NOX) family of enzymes generates the ROS. The regulation of NOX enzymes is complex, with many members of this family exhibiting complexity in terms of subunit composition, cellular location, and tissue-specific expression. Cells are continuously exposed to the ROS, which represent mutagens and are thought to be a major contributor to several diseases including cancer and aging process. Therefore, cellular ROS sensing and metabolism are firmly regulated by a variety of proteins involved in the redox mechanism. In this review, the roles of oxidative stress in PI3K/AKT/PTEN signaling are summarized with a focus on the links between the pathways and NOX protein in several diseases including cancer and aging.

Key words: PTEN, PI3K, AKT, ROS, PPAR, WRN, SIRT1, cell signaling

Oxidative stress results from an imbalance of an excess of oxidants relative to antioxidant capacity [1]. The oxidants derive from various oxidase enzymes, including xanthine oxidase, lipoxygenase, or nicotinamide adenine denucleotide phosphate (NADPH) oxidase (NOX). An endogenous reactive oxygen species (ROS) mainly comes from mitochondria during the process of oxidative phosphorylation to produce energy in the form of ATP [2]. The ROS are also produced by intracellular membrane oxidases. Inflammation is typical source of those ROS at the sites of tissues. Some ROS function as signaling molecules that mediate various cellular responses. Experimental evidences suggest that free radical production in cells is critical to intracellular signal transduction and regulate various physiological processes including proliferation, differentiation, apoptosis and migration [3, 4]. The NOX family is considered one of the sources of ROS which participates in the induction of

important signaling pathways. For example, it has been reported that p42/44 ERK activation is NOX dependent [5]. In addition, ROS modify the activity of several key enzymes for cell-dynamics, resulting in the reorganization of actin cytoskeleton, adhesion and cell migration [6]. On the other hand, ROS may react with cellular constituents including proteins, lipids, and DNA to generate an array of oxidative lesions. Protein and lipid oxidation by ROS is often proposed as a crucial determinant of health. These lesions may compromise genome stability which is critical for cellular homeostasis and progeny [7]. Increased levels of oxidative stress results in macromolecular damage and is implicated in various disease states such as atherosclerosis, diabetes, and cancer [8]. Accumulating evidence suggest that NOX-derived ROS may elevate the risk for genomic instability [9].

It is important for cells to neutralize ROS before they can damage cellular macromolecules. Therefore, cellular

*Correspondence should be addressed to: Dr. Satoru Matsuda, Department of Food Science and Nutrition, Nara Women's University, Kita-Uoya Nishimachi, Nara 630-8506, Japan. Email: smatsuda@cc.nara-wu.ac.jp

ROS metabolism is firmly regulated by a range of proteins involved in the redox mechanism. However, less is known about the initial regulation of signaling molecules for ROS production. NOX enzymes play central roles as they can stimulate other enzymatic sources of ROS. PI3K/AKT signaling pathway seems to play important roles in the NOX activation. It has been shown that the initial step in a NOX-activation requires PI3K by using a dominant negative construct. *PTEN* (phosphatase and tensin homolog deleted on chromosome 10) is a tumor suppressor gene that is frequently deleted or mutated in a variety of human cancers. It has been demonstrated that

up-regulation of PTEN causes modulation of PI3K/AKT signaling to reduce ROS generation in cells [10]. The PI3K/AKT/PTEN pathway may be involved in the mechanisms of several diseases. Accordingly, investigation of the interactions of redox- and non-redox-based signaling pathways could lead to better pharmacological control over the diseases. In this review, NOX family proteins and its roles in oxidative stress are summarized with a focus on the molecular links and interplay between NOX and the PI3K/AKT/PTEN signaling pathway.




Figure 1. Schematic structures of human PTEN and AKT1 protein. The predicted consensual domain structures for each protein are depicted. Note that the sizes of protein are modified for clarity. C2 domain= a protein structural domain involved in targeting proteins to cell membranes; PDZ= a common structural domain in signaling proteins (PSD95, Dlg, ZO-1, etc); PH domain= pleckstrin homology domain

Characteristics of PI3K/AKT/PTEN pathway and NOX proteins

The PIP3 is the major second messenger of the PI3K pathway that mediates receptor tyrosine kinase signaling to the proliferation and survival kinase AKT [11]. PTEN negatively regulates the activity of PI3K/AKT signaling through converting PIP3 to PIP2, by which PTEN exerts its tumor-suppressive effect. Increased levels of PIP3 at the membrane cause PH domain-containing protein kinases to co-localize, resulting in the kinases-mediated phosphorylation and activation [12]. For example, the activated AKT phosphorylates target proteins involved in

cell survival, cell cycling, angiogenesis, and metabolism [13]. PTEN acts as regulator of maintaining basal levels of PIP3 opposing to those signaling activation too much, as the increased proliferation, survival and motility are main cellular effects that contribute to tumorigenesis. Actually, dysregulation of the PI3K/AKT pathway has been found in many malignant cancers [14]. Human PTEN is a 53 kDa protein with homology to tensin and protein tyrosine phosphatases. Schematic structure of the PTEN protein is shown in Figure 1. The structure provides PTEN with its preference for acidic phospholipid substrates such as PIP3. Tumor suppressor p53 and peroxisome proliferator activated receptor γ (PPAR γ) can

transcriptionally upregulate *PTEN* expression [15, 16]. Interestingly, it is reported that rosemary extract represses *PTEN* expression in K562 leukemic culture cells [17]. Overexpression of *PTEN* induces growth suppression by promoting G1 arrest [18]. AKT activation leads to Hypoxia-inducible factor-1 (HIF-1) stabilization, whereas *PTEN* attenuates the hypoxia-mediated HIF-1 stabilization [19]. The HIF-1 is a critical element for the transcriptional regulation of genes important for adaptation to low oxygen conditions. The instability of mutant *PTEN* and the reduction of HIF1 degradation have been shown to involve protein interactions [20]. *PTEN* has been shown to act as a tumor suppressor whose function includes important roles in regulating oxidative stress, indicating a potential role in oxidative damage-associated diseases [21].

NOX family proteins are committed reactive oxygen species-generating enzymes, which also regulate redox-sensitive signaling pathways [22]. The NOX family has

been categorized into seven homologues (NOX1- NOX5 and DUOX1- DUOX2) depending on the membrane-associated flavin-proteins. NOXs are multimeric enzymes that consist of the cytosolic subunits p22phox, p40phox, p47phox, p67phox, and Rac1 [23] (Figure 2). Upon activation, all of them as well as the transmembrane glycoprotein gp91phox and p22phox translocate to the membrane [23]. NOX-related cytosolic proteins associate with integral membrane subunits to form the functional enzyme, which generates ROS. NOX catalyses a one-electron transfer to molecular oxygen to form the radical superoxide anion, which can be reduced to hydrogen peroxide. It has been reported that loss of fluid shear stress, ischemia, and inflammation cause plasma membrane depolarization via ATP-sensitive K channel closure, initiating a signaling cascade which leads to NOX activation and ROS production [24].




Figure 2. NOXs are trans-membrane protein and can form a complex with p22phox, Rac1, p40phox, p47phox and p67phox. NOXs are responsive to the extracellular fluid surrounding the inflammation and stress, and generate ROS. Note that some critical molecules have been omitted for clarity.

Roles of PI3K/ AKT /PTEN and NOX for the ROS production

Since wortmannin, a PI3K/AKT inhibitor, considerably inhibits the translocation of NOX subunits and reduces ROS production, in which the role of PI3K signaling might be involved [25, 26] (Figure 3). It has also been

reported that cell membrane depolarization, which is followed by PI3K/AKT and PKC activation causes NOX assembly and ROS production. The ROS production is abolished by the wortmannin as well as the PKC inhibitor H7 [24]. In addition, the combination of the wortmannin and H7 does not have a larger effect. In AKT1-null cells, the ROS production is reduced compared with wild-type

cells [27]. Although wortmannin prevents the NOX activation, wortmannin has no effect on the cell membrane depolarization. Activation of NOX is also diminished by dominant negative PI3K [28, 29]. Accordingly, AKT phosphorylation follows cell membrane depolarization, and it precedes the activation of NOX. In other words, PI3K/AKT and PKC activations are downstream of cell membrane depolarization but upstream of the NOX activation. It has also been shown

that ROS could play a critical role in pancreatic islet β -cell dysfunction through PTEN-dependent JNK activation and AKT inhibition [30]. These results indicate that PI3K/AKT and PKC are in the same pathway and are required for ROS production. However, the intermediate steps that link depolarization and the NOX activation are not well understood.




Figure 3. Schematic representation of PI3K/AKT/PTEN signaling, which modulate NOXs-derived ROS. NOXs sense hypoxia and can help to mediate activation of angiogenic pathway, while maintaining unregulated cell growth and apoptotic cell death. Examples of molecules known to act on the regulatory pathways are shown. Note that some critical pathways have been omitted for clarity.

ROS directly interact with signaling molecules to modulate the signaling in a variety of cellular processes such as proliferation and survival. Actually, the catalytic activity of PTEN is also modulated by ROS, and cellular PTEN phosphatase activity is inhibited by the oxidative stress [31]. The PTEN inactivation then causes an increase in cellular PIP3 levels, PIP3 accumulation occurs at the plasma membrane and activation of the downstream PIP3 target including AKT, indicating a part of functional role for elevated intracellular ROS. Furthermore, activated PI3K/AKT signaling causes increased expression of several genes for cell survival. Endogenous oxidant production in macrophages inactivates a fraction of the cellular PTEN, and this is associated with an oxidant-dependent activation of downstream signaling [32]. The phosphorylated PTEN inactivation and the consequent

AKT activation in cells are inhibited by antioxidant treatment. ROS levels are increased in the retinal pigment epithelium cells in association with phosphorylation and inactivation of PTEN [33]. Mitochondrial PTEN protein levels are increased by H₂O₂, although ROS does not affect the PTEN expression [34]. The increased PTEN in mitochondria may further promote ROS production in cells. However, exposure of the PTEN to H₂O₂ results in inactivation of PTEN [35]. Various cysteine mutants in the active site of PTEN have indicated that the essential Cys residue specifically forms a disulfide bond during oxidation [36]. Uncontrolled generation of ROS might contribute to the cell proliferation by inhibiting PTEN function.




Figure 4. The PI3K/AKT/ PTEN pathway and ROS may be involved in several diseases besides cancer and aging.

Link between PI3K/AKT/PTEN pathway and NOX protein involved in several Diseases

Intracellular signaling is initiated at low ROS concentration, whereas oxidative stress is induced at high ROS concentration. Therefore, imbalance of NOX activities could easily be the potential cause of some

diseases (Figure 4). With regard to functional relationships between NOX isoforms and pathogenesis, it is of particular interest to study whether they are involved in the development in several diseases, because overproduction of ROS has long been implicated as a risk factor. For example, multiple genetic and epigenetic alterations by oxidative stress are required for cellular

transformation and carcinogenesis. Inactivation of the PTEN is implicated in the tumorigenesis of prostate cancer, in which PIP3 and oxidants seem to be produced as carcinogens. It has been reported that causal relationship between NOX and cancer is also in Ras oncogene-induced cell transformation [37, 38]. NOX family of genes appears to be required for survival and growth of a subset of human cancer cells [37]. In addition, NOX-derived ROS are known to play a central role in each step of the metastatic cascade including invasion, adhesion, angiogenesis and proliferation. ROS attack guanine bases in DNA and form 8-hydroxydeoxyguanosine (8-OHdG), which can bind to thymidine rather than cytosine, based on which, the level of 8-OHdG is generally regarded as a biomarker of mutagenesis consequent to oxidative stress [38, 39]. Higher levels of 8-OHdG are noted in *Helicobacter pylori* infection in atrophic gastritis as well as in gastric cancer [38, 39]. They act at specific cellular microdomains through the activation of oncogenes and the inactivation of tumor suppressor proteins. The produced ROS surrounding the tumor environment will continue unregulated cell growth, progress, facilitate invasion, and make metastasis.

Defects in insulin secretion and decreased sensitivity of target tissues to insulin action are the key features of type 2 diabetes. Dysfunction of β -cell in pancreas is one of major characteristics in the pathogenesis of type 2 diabetes [40]. It has been shown that excessive generation of ROS is linked to the β -cell dysfunction. Oxidative stress has also been implicated in the pathogenesis of free fatty acids (FFAs)-induced β -cell dysfunction. The combination of obesity and type 2 diabetes is associated with elevated plasma FFAs that lead to apoptosis of β -cells through PTEN-dependent AKT inhibition, followed by reduced insulin expression and secretion [41]. Although NOX-generated ROS are the well characterized in mammalian cells, ROS are also generated by the oxidative metabolism of arachidonic acid that is released from the membrane phospholipids via the activity of cytosolic phospholipase A2 (PLA2) [42, 43]. The interaction between ROS and inflammation also plays an important role in some other disease pathogenesis. For example, chronic airway diseases such as asthma are linked to oxidative environmental factors and are associated with increased production of ROS [44]. Oxidative stress seems an important contributing factor to asthma, and therefore, antioxidant strategies may be useful in the treatment of asthma. It is not surprising that the contributions of NOX-derived ROS to various aspects of asthma development and progression are diverse. Thyroid cells contain antioxidants to protect cells from the ROS-mediated oxidative damage [45]. NOX4, a homolog of the NOX family, is an intracellular source of ROS in

the human thyroid gland. Papillary thyroid carcinoma, the most common thyroid cancer, is closely linked to the increased ROS production by NOX4 [46]. Imbalance of ROS may result in thyroid cell dysfunction and thyroid diseases, which provides us with valuable tools for a better understanding of pathophysiology of prevalent thyroid diseases. Furthermore, ROS regulates immune system and enhances atherosclerosis [47, 48].

Werner syndrome, which is characterized by the early onset of premature age-associated pathologies including elevated cancer incidence, is a rare autosomal disease [49]. Werner syndrome is caused by the lack of a functional nuclear RecQ-like DNA helicase, named WRN protein. The DNA damage, including oxidative DNA damage at telomeres, which plays a critical role in optimizing double strand breaks repair mechanisms due to its end-processing helicase and exonuclease activities. Studies indicate that WRN protein regulates HIF-1 activation by affecting the mitochondrial ROS production [50]. The depletion of WRN protein leads to increased HIF-1 complex stabilization and activation [50]. HIF-1 is implicated in the molecular mechanisms of ageing, whose activation in the absence of WRN involves the generation of mitochondrial ROS. Activation of PI3K/AKT pathway also leads to HIF-1 stabilization [51], whereas PTEN attenuates the HIF-1 stabilization [52]. Mutation of PTEN can destabilize the HIF-1. The deletion of AKT protects against the age-dependent defects, raising the possibility that modulation of the PTEN/AKT pathway could protect against premature aging syndromes. Down-regulation of PI3K/AKT pathway by PTEN expression may protect against age-dependent DNA damage and cancer, including the premature disorders observed in Werner syndrome. Vitamin C supplementation also rescues the shorter mean life span of the *WRN* mutant mice and reverses several abnormalities [53].

Discussion and Perspective

Induction of ROS is likely to play a key role in cell growth inhibitory responses. It is accepted that one of the symbols of aging is the accumulation of DNA damage caused by the ROS, which is known to induce genomic alterations such as point mutations and deletions, then inhibit tumor suppressor genes and/or activate oncogenes. It has been shown that tumor suppressor p53 also promotes ROS production and participates in the induction of apoptosis. Increasing ROS can also enhance insulin signaling to attenuate the development of insulin resistance. The enhanced ROS-dependent insulin signaling may be attributable to the oxidation and inhibition of PTEN. Given the importance of PTEN in regulating metabolism, there are many linking PTEN to diabetes and senescence. In addition, PTEN and PI3K/AKT are in the same

pathway and are supposed to be required for ROS production. As ROS can regulate PTEN, the role of PTEN in modulating ROS may form part of a feedback loop. However, this may paradoxically carry damaging consequences in the setting of persistent environmental stressors and during the process of aging. A better understanding of the complicated relationships between ROS and aging via NOX and PTEN/AKT pathway may provide clues to ROS-mediated pathogenesis-pathways for new drug discovery. Future appropriate experiments will determine which pathways are responsible for the redox imbalance.

Acknowledgments

This work was supported by grants-in-aid JSPS KAKENHI Grant Number 24240098 and 25560050 in Japan. In addition, this work was supported in part by the grant from Nakagawa Masashichi Shoten Co., Ltd. and SHIN-EI Pharmaceutical Co., Ltd.

Competing interests statement

The authors declare that they have no competing financial interests.

Reference

- [1] Salvemini D and Cuzzocrea S (2002). Oxidative stress in septic shock and disseminated intravascular coagulation. *Free Radic Biol Med*, 33:1173-1185
- [2] Ballard JW (2005). *Drosophila simulans* as a novel model for studying mitochondrial metabolism and aging. *Exp Gerontol*, 40: 763-773
- [3] Zhang Y, Du Y, Le W, Wang K, Kieffer N and Zhang J (2011). Redox control of the survival of healthy and diseased cells. *Antioxid Redox Signal*, 15: 2867-2908
- [4] Scatena R (2012). Mitochondria and cancer: a growing role in apoptosis, cancer cell metabolism and dedifferentiation. *Adv Exp Med Biol*, 942: 287-308
- [5] Wu RF, Ma Z, Myers DP and Terada LS (2007). HIV-1 Tat activates dual Nox pathways leading to independent activation of ERK and JNK MAP kinases. *J Biol Chem*, 282: 37412-37419
- [6] San Mart ín A and Griending KK (2010). Redox control of vascular smooth muscle migration. *Antioxid Redox Signal*, 12: 625-640
- [7] Weyemi U, Redon CE, Parekh PR, Dupuy C and Bonner WM (2013). NADPH Oxidases NOXs and DUOXs as putative targets for cancer therapy. *Anticancer Agents Med Chem*, 13: 502-514
- [8] Roberts CK and Sindhu KK (2009). Oxidative stress and metabolic syndrome. *Life Sci*, 84: 705-712
- [9] Leonarduzzi G, Sottero B, Testa G, Biasi F and Poli G (2011). New insights into redox-modulated cell signaling. *Curr Pharm Des*, 17: 3994-4006
- [10] Xu J, Tian W, Ma X, et al. (2011). The molecular mechanism underlying morphine-induced AKT activation: roles of protein phosphatases and reactive oxygen species. *Cell Biochem Biophys*, 61: 303-311
- [11] Maehama T, Taylor GS and Dixon JE (2001). PTEN and myotubularin: novel phosphoinositide phosphatases. *Annu Rev Biochem*, 70: 247-279
- [12] Howes AL, Arthur JF, Zhang T, et al. (2003). AKT-mediated cardiomyocyte survival pathways are compromised by G alpha q-induced phosphoinositide 4,5-bisphosphate depletion. *J Biol Chem*, 278: 40343-40351
- [13] Sheppard K, Kinross KM, Solomon B, Pearson RB and Phillips WA (2012). Targeting PI3 kinase/AKT/mTOR signaling in cancer. *Crit Rev Oncog*, 17: 9-95
- [14] Castaneda CA, Cortes-Funes H, Gomez HL and Ciruelos EM (2010). The phosphatidyl inositol 3-kinase/AKT signaling pathway in breast cancer. *Cancer Metastasis Rev*, 29: 751-759
- [15] Teresi RE, Shaiu CW, Chen CS, Chatterjee VK, Waite KA and Eng C (2006). Increased PTEN expression due to transcriptional activation of PPARgamma by Lovastatin and Rosiglitazone. *Int J Cancer*, 118: 2390-2398
- [16] Harikumar KB and Aggarwal BB (2008). Resveratrol: a multitargeted agent for age-associated chronic diseases. *Cell Cycle*, 7: 1020-1035
- [17] Yoshida H, Okumura N, Kitagishi Y, Nishimura Y and Matsuda S (2011). Ethanol extract of Rosemary repressed PTEN expression in K562 culture cells. *Int J appl Boil pharm Technol*, 2: 316-322
- [18] Weng LP, Brown JL and Eng C (2001). PTEN coordinates G(1) arrest by down-regulating cyclin D1 via its protein phosphatase activity and up-regulating p27 via its lipid phosphatase activity in a breast cancer model. *Hum Mol Genet*, 10: 599-604
- [19] Carver DJ, Gaston B, Deronde K and Palmer LA (2007). AKT-mediated activation of HIF-1 in pulmonary vascular endothelial cells by S-nitrosoglutathione. *Am J Respir Cell Mol Biol*, 37: 255-263
- [20] Li YM, Zhou BP, Deng J, Pan Y, Hay N and Hung MC (2005). A hypoxia-independent hypoxia-inducible factor-1 activation pathway induced by phosphatidylinositol-3 kinase/AKT in HER2 overexpressing cells. *Cancer Res*, 65: 3257-3263
- [21] Kitagishi Y and Matsuda S (2013). Redox regulation of tumor suppressor PTEN in cancer and aging (Review). *Int J Mol Med*, 31: 511-515
- [22] Maghzal GJ, Krause KH, Stocker R and Jaquet V (2012). Detection of reactive oxygen species derived from the family of NOX NADPH oxidases. *Free Radic Biol Med*, 53: 1903-1918
- [23] El-Benna J, Dang PM, Gougerot-Pocidallo MA, Marie JC and Braut-Boucher F (2009). p47phox, the phagocyte NADPH oxidase/NOX2 organizer: structure, phosphorylation and implication in diseases. *Exp Mol Med*, 41: 217-225
- [24] Chatterjee S, Browning EA, Hong N, DeBolt K, Sorokina EM, Liu W, Birnbaum MJ and Fisher AB (2012). Membrane depolarization is the trigger for PI3K/Akt activation and leads to the generation of ROS.

- Am J Physiol Heart Circ Physiol, 302: H105-H114
- [25] Ren Z, Raucci FJ Jr, Browe DM and Baumgarten CM (2008). Regulation of swelling-activated Cl(-) current by angiotensin II signalling and NADPH oxidase in rabbit ventricle. *Cardiovasc Res*, 77: 73-80
- [26] Ngkelo A, Meja K, Yeadon M, Adcock I and Kirkham PA (2012). LPS induced inflammatory responses in human peripheral blood mononuclear cells is mediated through NOX4 and G α dependent PI-3kinase signalling. *J Inflamm (Lond)*, 9:1. doi: 10.1186/1476-9255-9-1.
- [27] Jee HJ, Kim HJ, Kim AJ, Bae YS, Bae SS and Yun J (2009). UV light induces premature senescence in Akt1-null mouse embryonic fibroblasts by increasing intracellular levels of ROS. *Biochem Biophys Res Commun*, 383: 358-362
- [28] B äumer AT, Ten Freyhaus H, Sauer H, Wartenberg M, Kappert K, Schnabel P, Konkol C, Hescheler J, Vantler M and Rosenkranz S (2008). Phosphatidylinositol 3-kinase-dependent membrane recruitment of Rac-1 and p47phox is critical for alpha-platelet-derived growth factor receptor-induced production of reactive oxygen species. *J Biol Chem* 283: 7864-7876
- [29] Lee SB, Bae IH, Bae YS and Um HD (2006). Link between mitochondria and NADPH oxidase 1 isozyme for the sustained production of reactive oxygen species and cell death. *J Biol Chem*, 281: 36228-36235
- [30] Yuan H, Lu Y, Huang X, He Q, Man Y, Zhou Y, Wang S and Li J (2010). Suppression of NADPH oxidase 2 substantially restores glucose-induced dysfunction of pancreatic NIT-1 cells. *FEBS J*, 277: 5061-5071
- [31] Barbieri SS, Ruggiero L, Tremoli E and Weksler BB (2008). Suppressing PTEN activity by tobacco smoke plus interleukin-1beta modulates dissociation of VE-cadherin/beta-catenin complexes in endothelium. *Arterioscler Thromb Vasc Biol*, 28: 732-738
- [32] Leslie NR, Bennett D, Lindsay YE, Stewart H, Gray A and Downes CP (2003). Redox regulation of PI 3-kinase signalling via inactivation of PTEN. *EMBO J*, 22: 5501-5510
- [33] Kim JW, Kang KH, Burrola P, MAKTW and Lemke G (2008). Retinal degeneration triggered by inactivation of PTEN in the retinal pigment epithelium. *Genes Dev*, 22: 3147-3157
- [34] Zu L, Zheng X, Wang B, et al. (2011). Ischemic preconditioning attenuates mitochondrial localization of PTEN induced by ischemia-reperfusion. *Am J Physiol Heart Circ Physiol*, 300: H2177-H2186
- [35] Lee SR, Yang KS, Kwon J, Lee C, Jeong W and Rhee SG (2002). Reversible inactivation of the tumor suppressor PTEN by H₂O₂. *J Biol Chem*, 277: 20336-20342
- [36] Matsuda M, Takeshita K, Kurokawa T, et al. (2011). Crystal structure of the cytoplasmic phosphatase and tensin homolog (PTEN)-like region of *Ciona intestinalis* voltage-sensing phosphatase provides insight into substrate specificity and redox regulation of the phosphoinositide phosphatase activity. *J Biol Chem*, 286: 23368-23377
- [37] Kamata T (2009). Roles of Nox1 and other Nox isoforms in cancer development. *Cancer Sci*, 100: 1382-1388
- [38] Shinohara M, Shang WH, Kubodera M, Harada S, Mitsushita J, Kato M, Miyazaki H, Sumimoto H and Kamata T (2007). Nox1 redox signaling mediates oncogenic Ras-induced disruption of stress fibers and focal adhesions by down-regulating Rho. *J Biol Chem*, 282: 17640-17648
- [39] Ock CY, Kim EH, Choi DJ, Lee HJ, Hahn KB and Chung MH (2012). 8-Hydroxydeoxyguanosine: not mere biomarker for oxidative stress, but remedy for oxidative stress-implicated gastrointestinal diseases. *World J Gastroenterol*, 18: 302-308
- [40] Cefalu WT (2006). Animal models of type 2 diabetes: clinical presentation and pathophysiological relevance to the human condition. *ILAR J*, 47: 186-198
- [41] Yuan H, Zhang X, Huang X, Lu Y, Tang W, Man Y, Wang S, Xi J and Li J (2010). NADPH oxidase 2-derived reactive oxygen species mediate FFAs-induced dysfunction and apoptosis of β -cells via JNK, p38 MAPK and p53 pathways. *PLoS One*, 5: e15726
- [42] Muralikrishna Adibhatla R and Hatcher JF (2006). Phospholipase A2, reactive oxygen species, and lipid peroxidation in cerebral ischemia. *Free Radic Biol Med*, 40: 376-387
- [43] Adibhatla RM and Hatcher JF (2008). Phospholipase A(2), reactive oxygen species, and lipid peroxidation in CNS pathologies. *BMB Rep*, 41: 560-567
- [44] Wood LG, Wark PA and Garg ML (2010). Antioxidant and anti-inflammatory effects of resveratrol in airway disease. *Antioxid Redox Signal*, 13: 1535-1548
- [45] Chiu-Ugalde J, Wirth EK, Klein MO, Sapin R, Fradejas-Villar N, Renko K, Schomburg L, Kährle J and Schweizer U (2012). Thyroid function is maintained despite increased oxidative stress in mice lacking selenoprotein biosynthesis in thyroid epithelial cells. *Antioxid Redox Signal*, 17: 902-913
- [46] Weyemi U, Caillou B, Talbot M, Ameziane-El-Hassani R, Lacroix L, Lagent-Chevallier O, Al Ghuzlan A, Roos D, Bidart JM, Virion A, Schlumberger M and Dupuy C (2010). Intracellular expression of reactive oxygen species-generating NADPH oxidase NOX4 in normal and cancer thyroid tissues. *Endocr Relat Cancer*, 17: 27-37
- [47] Ishiyama J, Taguchi R, Yamamoto A and Murakami K (2010). Palmitic acid enhances lectin-like oxidized LDL receptor (LOX-1) expression and promotes uptake of oxidized LDL in macrophage cells. *Atherosclerosis*, 209: 118-124
- [48] Laddha NC, Dwivedi M, Mansuri MS, Gani AR, Ansarullah M, Ramachandran AV, Dalai S and Begum R (2013). Vitiligo: interplay between oxidative stress and immune system. *Exp Dermatol*, 22: 245-250
- [49] Wirtenberger M, Frank B, Hemminki K, Klaes R, Schmutzler RK, Wappenschmidt B, Meindl A, Kiechle M, Arnold N, Weber BH, Niederacher D, Bartram CR and Burwinkel B (2006). Interaction of Werner and Bloom syndrome genes with p53 in familial breast cancer. *Carcinogenesis*, 27: 1655-1660
- [50] Labbé A, Lafleur VN, Patten DA, Robitaille GA, Garand C, Lamallice L, Lebel M and Richard DE (2012). The Werner syndrome gene product (WRN): a repressor of

- hypoxia-inducible factor-1 activity. *Exp Cell Res*, 318: 1620-1632
- [51] Shafee N, Kaluz S, Ru N and Stanbridge EJ (2009). PI3K/Akt activity has variable cell-specific effects on expression of HIF target genes, CA9 and VEGF, in human cancer cell lines. *Cancer Lett*, 282: 109-115
- [52] Zundel W, Schindler C, Haas-Kogan D, Koong A, Kaper F, Chen E, Gottschalk AR, Ryan HE, Johnson RS, Jefferson AB, Stokoe D and Giaccia AJ (2000). Loss of PTEN facilitates HIF-1-mediated gene expression. *Genes Dev*, 14: 391-396
- [53] Massip L, Garand C, Paquet ER, et al. (2010). Vitamin C restores healthy aging in a mouse model for Werner syndrome. *FASEB J*, 24: 158-172