

# Evidence that Increased Acetaminophen use in Genetically Vulnerable Children Appears to be a Major Cause of the Epidemics of Autism, Attention Deficit with Hyperactivity, and Asthma

William Shaw, PhD<sup>a</sup>

©2013, William Shaw, PhD  
Journal Compilation ©2013, AARM  
DOI 10.14200/jrm.2013.2.0101

## ABSTRACT

It appears that the marked increase in the rate of autism, asthma, and attention deficit with hyperactivity throughout much of the world may be largely caused by the marked increase in the use of acetaminophen in genetically and/or metabolically susceptible children, and the use of acetaminophen by pregnant women. Toxicity of acetaminophen may cause autism by overloading the defective sulfation pathway catalyzed by phenolsulfotransferase, which is deficient in autism, leading to overproduction of the toxic metabolite *N*-acetyl-*p*-benzoquinone imine (NAPQI). Increased levels of NAPQI reduce the ability to detoxify a host of toxic chemicals in the environment, increasing oxidative stress, which leads to protein, lipid, and nucleic acid damage from free radicals. Epidemiological evidence also supports the association of increased acetaminophen usage with autism, cough, and attention deficit with hyperactivity. The marked increases in the incidences of autism, asthma, and attention deficit disorder in the United States coincide with the replacement of aspirin by acetaminophen in the 1980s. The characteristic loss of Purkinje cells in the brains of people with autism is consistent with depletion of brain glutathione due to excess acetaminophen usage, which leads to premature brain Purkinje cell death. The anomalous hair mercury concentrations of children with autism are consistent with exposure of growing hair proteins to NAPQI derived from acetaminophen, which competitively inhibits the reaction of mercury with hair sulfhydryl groups. Finally, large-scale faulty production of acetaminophen products, such that the labeled values were exceeded by the true concentrations, in addition to contamination with bacteria and tribromoanisole, may have greatly increased the chances of children receiving overdosages of acetaminophen and potential toxins for perhaps as long as a decade.

**Keywords:** Acetaminophen; Asthma; Attention deficit with hyperactivity; Autism; Glutathione; *N*-acetyl-*p*-benzoquinone imine (NAPQI); Phenolsulfotransferase; Tribromoanisole

<sup>a</sup> Director, The Great Plains Laboratory, Inc., 11813 West 77th Street, Lenexa, KS 66214,  
Tel.: 913-341-8949, E-mail: wshaw@gpl4u.com

## INTRODUCTION

One of the puzzling aspects of autism is the marked increase in the incidence of autism that began in the United States in the early 1980s and has appeared to increase continuously since then. The highest incidence of autism has been reported to be South Korea, where the incidence is now reported to be one in 38 boys.<sup>1</sup> Increased incidence of autism due to more effective diagnosis was disproved in the study of Irva Hertz-Picciotto who showed that perhaps 12% of the increased autism diagnoses could be attributed to improved diagnosis.<sup>2</sup> A wide range of environmental factors has been associated with increased autism incidence, including pesticides, chemicals, phthalates, polychlorinated biphenyls, solvents, heavy metals or other pollutants.<sup>3</sup> Although toxic chemicals are undoubtedly not beneficial for the health of any person, is there any information that indicates that a toxic avalanche of chemicals inundated the United States in the early 1980s? Indeed a wealth of knowledge about environmental chemicals has led to marked reductions in exposure to chemicals such as lead and dichlorodiphenyltrichloroethane (DDT) in the United States over the past 50 years. For example, acceptable safe limits for levels of lead in the blood have decreased from 60 µg/dL in 1960 to <5 µg/dL in 2010.<sup>4</sup>

Making a connection between disease appearance and causative agent is important. Clinical studies, epidemiological studies and post-market pharmacovigilance are of utmost importance in recognizing signals and drug-induced side effects. One of the most notable cases of serious adverse effects caused by a pharmaceutical agent was the terrible developmental epidemic of the birth of children with seal-like arms and legs (phocomelia) that was linked to the maternal use of the sedative thalidomide 20–35 days after conception.<sup>5</sup> What would have happened if the thalidomide connection had never been made? One of the difficulties with chemical studies of autism associations is that most chemicals are used worldwide, making it difficult to find a “clean” environment where autism might be less prevalent. One of the clues that led to the discovery of thalidomide as the causative agent of deformed limbs was that it was much more commonly used in Europe than in the United States. Countries with the

greatest use of thalidomide by pregnant women during pregnancy were those with the highest incidence of deformed babies. If there was a geographic region in the world in which the incidence of autism was much lower than that in the United States, a comparison of medical or dietary differences might provide a significant clue to the major cause of autism.

Such a country is Cuba. The highest estimate of the total incidence of autism in Cuba is 185 cases out of a total population of 11,000,000 (0.00168% of the population) compared with an estimate of as high as 1.5 million in a total United States population of 300 million (0.50%).<sup>6,7</sup> The percentage of the population with autism in the United States is thus 298 times higher than in Cuba. Cuba is much more economically challenged than the United States, with the per capita income of Cuba approximately eight times lower than that in the United States. Despite the economic challenges presented to the communist government of Cuba, basic healthcare is readily available and there are a large number of physicians trained in 14 different medical schools. Unlike the United States, where vaccines are optional in many states, vaccines are compulsory in Cuba and Cuba has one of the most highly vaccinated populations in the world against a wide variety of infectious agents.<sup>8</sup> For example, the vaccination rate for measles was reported to be 99.7%. The association of autism with various vaccines has had a very controversial history with inflamed passions on both sides of the debate and will not be examined here.

However, a topic much less frequently addressed in association with autism is the therapies that are given *in conjunction with vaccines*. The practice of prescribing acetaminophen as a prophylactic fever preventative is widespread in the United States but is very uncommon in Cuba (personal communications, Dr Olympio Rodriguez Santos MD, MSc, Allergist, Camaguey, Cuba). In the United States, some physicians have started to advise parents to begin to take acetaminophen prophylactically daily 5 days prior to childhood vaccines; some children on such prophylactic treatment had an autistic regression that began prior to vaccination (personal communication, Kerry Scott Lane MD, Anesthesiologist, West Palm Beach, Florida, USA). In Cuba, acetaminophen is

not approved as an over-the-counter (OTC) product, however, it has been available as an OTC product since 1959 in the United States. Furthermore, in Cuba, prophylactic use of antipyretic drugs is not the standard medical treatment for vaccine-related fever (personal communications with Dr Olympio Rodriguez Santos). If high fever continues after vaccination in Cuba for more than 2 days, the parents are advised to visit the physician's office where the drug metamizole is most commonly prescribed. Prescription of acetaminophen in such cases is rare. Metamizole is used in many countries throughout the world but is banned in the United States and some other countries because of a rare association with agranulocytosis.<sup>9</sup>

## COULD THE USE OF CERTAIN ANTIPYRETIC DRUGS, ESPECIALLY IN CONJUNCTION WITH VACCINES, BE A CAUSE OF AUTISM?

Torres was the first to ask if the suppression of fever by antipyretic drugs commonly used at the

time of vaccination might cause the severe immune abnormalities that are prevalent in autism.<sup>10</sup> Schultz *et al.* were even more specific when indicating in a series of articles that biochemical and immune disorders caused by increased use of the common drug acetaminophen may have caused the autism epidemic (Figure 1).<sup>11–14</sup>

Acetaminophen is also termed paracetamol and *N*-acetyl-*p*-aminophenol (AAP or APAP). More than 70% of the population in western countries has taken acetaminophen at least once, and a relevant percentage takes the drug chronically as a mild pain reliever and antipyretic.<sup>15</sup> Acetaminophen is used to treat pain and fever and it has become one of the most popular OTC non-narcotic analgesic agents. For example, this compound has been taken at least once by >85% of children under the age of 91 months in the UK.<sup>15</sup> In the US, approximately 79% of the general population regularly takes acetaminophen, including more than 35% of pregnant women.<sup>15</sup> Acetaminophen has grown in popularity in large part due to its reputation for safety. For generations, Tylenol® (a popular brand of acetaminophen) advertisements have painted it as “the pain reliever hospitals use most.” Acetaminophen is in >600 OTC




Figure 1: Increases in rates of asthma and autism with changes in acetaminophen usage. Modified from Reference 14

and prescription products, from headache and cold remedies to cough syrups and sleep aids.<sup>16</sup>

The study by Schultz *et al.* was the first to specifically link increased acetaminophen use to increased autism.<sup>11</sup> This study included a graph similar to Figure 1 temporally relating increased autism incidence in California with increased acetaminophen use in the United States and decreased acetaminophen use with decreased rate of autism in California. In addition, similar increases in the rates of asthma correlated with the usage of acetaminophen were also noted by Becker and Schultz.<sup>14</sup> They noted that the rates of autism incidence and asthma stopped increasing in the months following two attempted extortion events in which acetaminophen was deliberately laced with cyanide.<sup>14</sup> The changes in the incidence of these very different diseases at exactly the time acetaminophen use dropped for a significant period of time is remarkable and may indicate the same factor as causing the two diseases. In addition, the autism paper by Schultz *et al.* included the results of an online survey of parents who had given their child the combined measles, mumps, rubella (MMR) vaccine, which revealed that children with autism had more adverse reactions to the MMR vaccine and were more likely to have been given acetaminophen than ibuprofen for those reactions.<sup>11</sup> Compared with controls, children aged 1–5 years with autism were eight times more likely to have become unwell after the MMR vaccine, and were six times more likely to have taken acetaminophen. Children with autism who regressed in development were four times more likely to have taken acetaminophen after the vaccine. Illnesses concurrent with the MMR vaccine were nine times more likely in autistic children when all cases were considered, and 17 times more likely after limiting cases to children who regressed. There was no increased incidence of autism associated with ibuprofen use.

The incidence of attention deficit with hyperactivity over the last 50 years follows patterns similar to those of autism and asthma, although the data for attention deficit-hyperactivity disorder (ADHD) are not available to the same depth as the data for autism and asthma. Before 1970, the diagnosis of ADHD was relatively rare for schoolchildren and almost nonexistent for adolescents and adults. Between 1980 and 2007, there was an almost 8-fold increase of ADHD prevalence in the United States compared with rates of 40 years ago.<sup>17</sup> Prevalence

of ADHD in American schoolchildren was 1% in the 1970s, 3–5% in the 1980s, and 4–5% in the mid-to-late 1990s.<sup>18–24</sup> A study of hospital discharge rates for ADHD between 1989 and 2000 found a 381% increase over the study period.<sup>25</sup>

Use of acetaminophen dramatically increased in the United States in the 1980s due to a concern over an association of aspirin with Reye's syndrome, although a number of critics reject this hypothesis.<sup>26–28</sup> For example, the current recommendations for the management of children with Kawasaki disease include treatment with high-dose aspirin in the acute phase, and low-dose aspirin during the period of thrombocytosis. For those with residual coronary problems, low-dose aspirin is often given over an even longer term. In Japan alone, up to 200,000 children have received aspirin for Kawasaki disease. Interestingly, only *one* case of Reye's syndrome associated with Kawasaki disease has ever been reported, and only in the Japanese literature, giving an incidence of 0.005%.<sup>26</sup> In addition, retrospective reevaluation of patients with a diagnosis of Reye's syndrome who survived has revealed that many, if not most, had an underlying inborn error of metabolism (IEM).<sup>28</sup> Many of these IEMs had not even been described when the diagnosis of Reye's syndrome was made. Inborn errors that may mimic Reye's syndrome include fatty-acid oxidation defects, amino and organic acidopathies, urea-cycle defects, and disorders of carbohydrate metabolism.<sup>28</sup> Future discovery of other IEMs may ultimately explain even more of these cases. Additional etiologies that may mimic Reye's syndrome include viral infections, neuromuscular diseases, adverse drug reactions, and exposure to toxic chemicals and plants that cause hepatocellular damage and encephalopathy.<sup>28</sup> Diagnostic methods such as GC/MS became more widely available in the 1980s and later so that the patients with IEMs were diagnosed with an IEM instead of Reye's syndrome. The main cause of Reye's syndrome appears to be the accumulation of nonesterified fatty acids and lysolipids that have a high detergent activity and thus denature *all* proteins.<sup>29</sup>

It is interesting that Cuba, which has a lower autism rate than the United States, allows the use of acetaminophen only by prescription, therefore, the use of acetaminophen in Cuba is only a minuscule fraction of acetaminophen use in the United States and many other countries throughout the world. When



**Table 1:**

Harmful effects of acetaminophen or its metabolites	References
Neurotoxic effects on rat brain neurons both <i>in vitro</i> and <i>in vivo</i>	13, 15
Maternal use during pregnancy is associated with teratogenic defects in testicular function and gastrointestinal tract	30, 31
Very toxic metabolites formed from or induced by acetaminophen (NAPQI and peroxynitrite) that can cause oxidative damage to proteins, nucleic acids, amino acids, and lipids, causes increased mitochondrial and cellular damage and death	15, 32–35
Severe immune abnormalities at doses that do not damage the liver, including depression of specific antibody production and reduction of specific B-cells	36
Depresses the immune response to vaccination in humans and animals	37, 38
Depletion of glutathione by acetaminophen causes severe metabolic acidosis	39, 40
The leading cause of liver failure in the United States	14, 41, 42
A major cause of side effects that lead to emergency room visits in the United States (56,000 per year in the United States)	41, 42
Associated with increased rates of certain blood cancers	43
Acetaminophen prenatally or postnatally is associated with increased asthma	14, 44–46

NAPQI, *N*-acetyl-*p*-benzoquinone imine.

acetaminophen use was limited by a prescription requirement in the United States, the rate of autism was a small fraction of current rates of autism.

Unlike thalidomide, which was once promoted for its extreme safety prior to the discovery of its teratogenicity, acetaminophen has a long history of serious side effects associated with its use (Table 1).<sup>47</sup>

Acetaminophen produces neurotoxic effects on rat brain neurons both *in vitro* and *in vivo*, its use during pregnancy is associated with teratogenic defects in testicular function and the gastrointestinal tract, and there is increased incidence of asthma in maternally exposed and postnatally exposed children.<sup>13–15, 30, 31, 44–46</sup> Acetaminophen is converted to the very toxic metabolite *N*-acetyl-*p*-benzoquinone imine (NAPQI; Figure 2), which can cause oxidative damage to proteins, nucleic acids, amino acids, and lipids, in addition to increased mitochondrial and cellular damage and death.<sup>32–35</sup>


Acetaminophen also causes severe immune abnormalities at doses that do not damage the liver, depresses the immune response to vaccination, can cause severe metabolic acidosis when glutathione (GSH) is depleted, is the leading cause of liver failure and death in the United States, is associated with increased rates of certain blood cancers, and results in tens of thousands of visits to the emergency room and hospitalizations in the United States.<sup>14, 36–43</sup> A PubMed search of the scientific

literature indicated the presence of 2685 articles regarding acetaminophen toxicity.

An article with the title, “Did acetaminophen cause the autism epidemic?” was more pointed.<sup>48</sup> The rest of this article will deal with the known toxicity of acetaminophen and how other known anatomical, immunologic, biochemical, and infectious aspects of autism can be related to the effects of acetaminophen.

The metabolism of acetaminophen is shown in Figure 2. There are four major pathways for its detoxification. Acetaminophen can be converted to acetaminophen sulfate by phenol sulfotransferase (PST). It can also be converted to a glucuronide or deacetylated to a phenol. In addition, it can be converted to its extremely toxic metabolite NAPQI by the cytochrome P450 (CYP)2E1 enzyme. Several factors that increase the induction of CYP2E1, including smoking, obesity, vinyl chloride, and exposure to trichloroethylene are associated with increased incidence of autism. In addition, acetaminophen exposure itself induces increased CYP2E1, thus increasing the amount of NAPQI formed with each exposure to the drug.<sup>49–53</sup> Posadas *et al.* found that acetaminophen exposure to rats resulted in a concentration-related increase in CYP2E1 in rat brains.<sup>15</sup>

Thus, the prophylactic use of acetaminophen for days before vaccination and for multiple vaccinations would likely greatly increase the conversion of acetaminophen to its extremely toxic NAPQI metabolite. Glucuronidation is commonly present



**Figure 2: Metabolism of acetaminophen (paracetamol).**

CYP 450, cytochrome P450; NAPQI, *N*-acetyl-*p*-benzoquinone imine; PST, phenolsulfotransferase.

at low capacity in the fetus, newborns, and young infants, such that exposure to acetaminophen at these times leads to greater metabolism by other pathways.<sup>54</sup> Acetaminophen can also be deacetylated to form *p*-aminophenol, which can also be sulfated or converted to an active cannabinoid substance by being conjugated with arachidonic acid to form the conjugate termed AM404.<sup>55</sup> *p*-Aminophenol may also be detoxified by PST.

Acetaminophen produces analgesia by the activation of the brain endocannabinoid receptor CB1 by AM404.<sup>12, 13</sup> If the sulfation pathway is defective, as has been shown in autism, and/or there is impaired glucuronidation, acetaminophen will be increasingly converted to the alternative metabolic routes, increasing production of its more toxic compounds NAPQI and AM404.<sup>56, 57</sup>

In addition to liver toxicity caused by acetaminophen, individuals *without* liver toxicity may have severe metabolic acidosis if they are poorly nourished due to illness or dietary insufficiency.<sup>39, 40</sup> In

each of these cases, extremely high levels of pyroglutamic acid (also termed 5-oxoproline) are found in both the urine and blood serum. The connection between the elevation of this organic acid and acetaminophen use was first reported by Pitt *et al.* who suggested that acetaminophen ingestion depletes intracellular GSH stores, which then causes loss of feedback inhibition of  $\gamma$ -glutamylcysteine synthetase activity (Figures 3a and 3b).<sup>40</sup> This increases production of  $\gamma$ -glutamylcysteine, which is partially converted to pyroglutamic acid.

CYP enzymes are a superfamily of hemoproteins that carry out oxidative metabolism of many endogenous and foreign chemicals.<sup>54</sup> CYP2E1 is the principal CYP responsible for the metabolism of ethanol and is considered to be a major component of the microsomal ethanol-oxidizing system. Among xenobiotics metabolized by CYP2E1 are acetaldehyde, acetaminophen, acrylamide, aniline, benzene, butanol, carbon tetrachloride, diethylether, dimethyl sulfoxide, ethyl carbamate, ethylene chloride, halothane, glycerol, ethylene




Figure 3a: Metabolism of GSH in the absence of a toxic acetaminophen load.

glycol, *N*-nitrosodimethylamine, 4-nitrophenol, pyrazole, pyridine, and vinyl chloride.<sup>35</sup> Many of these chemicals are known toxins, established chemical carcinogens, or suspected carcinogens.

When *cyp2e1* knockout mice that lack the ability to produce the CYP2E1 enzyme were challenged with acetaminophen, they were found to be considerably *less sensitive* to its hepatotoxic effects than




Figure 3b: Metabolism of GSH after exposure to high doses of acetaminophen.

wild-type animals, indicating that this CYP is the principal enzyme responsible for the metabolic conversion of the drug acetaminophen to its active hepatotoxic metabolite NAPQI (Figure 2).<sup>35</sup> During fasting and diabetic ketosis, serum acetone, acetol, and 1,2-propanediol are elevated. CYP2E1 is concomitantly induced due to protein stabilization by acetone.<sup>35</sup> Acetone is primarily oxidized to acetol by CYP2E1. As fasting increases ketone production with a concomitant increase in CYP2E1 activity, this in turn increases the production of NAPQI and decreases amino acid substrates for GSH production.<sup>35</sup> Administering acetaminophen is likely to be much more toxic under fasting conditions, such as when a child has an illness that decreases appetite. Children who are sick and fasting and are administered vaccines with prophylactic acetaminophen are much more likely to suffer acetaminophen toxicity.

An examination of the importance of GSH and its biosynthesis is important to an understanding of acetaminophen toxicity. GSH is a tripeptide composed of the amino acids glutamate, cysteine and glycine (Figure 3a). It is present in virtually all aerobic cells at millimolar concentrations where it takes part in numerous fundamental processes. It is a very important antioxidant, participates in detoxification of certain drugs, toxic environmental chemicals, protects against lipid peroxidation and electrophiles, has antiviral effects, is involved in the biosynthesis of DNA, proteins, and leukotrienes, cell proliferation, apoptosis, neurotransmission, and neuromodulation.<sup>58</sup> Decreased levels of GSH are found in several diseases such as liver cirrhosis, pulmonary disease, gastrointestinal or pancreatic inflammation, diabetes, HIV infection, and neurodegenerative diseases.<sup>58</sup> In the normal physiological state (Figure 3a), GSH is produced by the condensation of the amino acids glutamate and cysteine to form the dipeptide  $\gamma$ -glutamylcysteine, which is catalyzed by the enzyme,  $\gamma$ -glutamyl cysteine synthetase.<sup>58</sup> The dipeptide then condenses with the amino acid glycine to form the tripeptide GSH which is catalyzed by the enzyme GSH synthetase. The end product, GSH, exhibits negative feedback inhibition of  $\gamma$ -glutamylsynthetase to prevent the overproduction of GSH. If the amino acid glycine is present at high concentrations,  $\gamma$ -glutamylcysteine is converted in small amounts to pyroglutamic acid that can be recycled to form glutamate.

If GSH is severely depleted by the toxic metabolite of acetaminophen, NAPQI (Figure 3b), and/or there is inadequate glycine to produce GSH due to illness or nutritional deficiency, there is a lack of negative feedback of GSH on  $\gamma$ -glutamylsynthetase.

This severe depletion of GSH results in the synthesis of large amounts of  $\gamma$ -glutamylcysteine, which is increasingly converted to pyroglutamate when glycine is depleted, leading to metabolic acidosis.

As organic acid testing to determine pyroglutamate is usually performed only at specialized pediatric hospitals with specialized biochemical genetics services, the diagnosis of this metabolic disorder in patients may frequently be missed. In addition, NAPQI also deactivates some of the enzymes that recycle GSH, such as GSH peroxidase.<sup>33</sup> The depletion of GSH diminishes the ability of the body to detoxify toxic chemicals. As of 2012, there were 170 articles that indicated an association between toxic chemical exposure and autism.<sup>3</sup> The depletion of GSH will lead to enhanced toxicity of a large number of toxic chemicals. For example, Adams *et al.* found that variations in the severity of autism measurements could be explained, in part, by regression analyses of urinary excretion of toxic metals before and after DMSA, the chelating agent dimercaptosuccinic acid, and the level of red blood cell glutathione.<sup>59</sup> Thus, partial depletion of GSH by moderate increases in NAPQI could lead to enhanced toxicity of heavy metals and perhaps many other toxic chemicals.

Depletion of GSH as a consequence of acetaminophen toxicity to the liver has attracted the most attention in the medical scientific community, as it can frequently be fatal or require a liver transplant or emergency treatment to prevent liver failure (the liver is the organ with the greatest concentration of GSH). However, acetaminophen toxicity has been implicated in a wide range of other disorders in humans and/or experimental animals including cancer, birth defects, asthma, allergies, and brain toxicity (Table 1).

Concentrations of *p*-aminophenol (which is converted to the cannabinoid substance AM404; Figure 2) from 1 to 100  $\mu\text{g/mL}$  produced significant loss of mouse cortical neuron viability at 24 h compared with the controls.<sup>11</sup> The naturally occurring endocannabinoid anandamide also caused similar 24 h



loss of cell viability in developing mouse cortical neurons at concentrations ranging from 1 to 100 µg/mL, indicating the mechanism of cell death could be mediated through the cannabinoid receptors. Defective glucuronidation would also increase the conversion of acetaminophen to its more toxic metabolites, NAPQI and AM 404. Such defects in glucuronidation are common in the developing fetus and in newborns.<sup>54</sup>

Posadas *et al.* found that acetaminophen causes concentration-dependent neuronal death *in vitro* at concentrations that are reached in human plasma during acetaminophen overdose, and are reached in the cerebrospinal fluid of rats for 3 h following doses that are below those required to induce acute hepatic failure in rats.<sup>15</sup> Acetaminophen also increases both neuronal CYP2E1 enzymatic activity and protein levels, as determined by western blot, leading to neuronal death through mitochondrial-mediated mechanisms that involve cytochrome c release and caspase 3 activation. In addition, *in vivo* experiments show that acetaminophen injection induces neuronal death in the rat cortex. Posadas *et al.* established a direct neurotoxic action by acetaminophen both *in vitro* and *in vivo* in rats at doses below those required to produce hepatotoxicity and suggested that this neurotoxicity might be involved in the general toxic syndrome observed during patient acetaminophen overdose and, possibly, when acetaminophen doses in the upper dosing schedule are used, especially if other risk factors (moderate alcohol drinking, fasting, nutritional impairment) are present.<sup>15</sup>

## PURKINJE CELL ABNORMALITIES, AUTISM, AND GSH DEPLETION

Ritvo *et al.* were the first to report abnormalities of Purkinje cells in the cerebella of people with autism.<sup>60</sup> These cells are some of the largest neurons in the human brain, with an intricately elaborate dendritic arbor, characterized by a large number of dendritic spines; these neurons utilize γ-aminobutyric acid as their neurotransmitter and send inhibitory projections to the deep cerebellar nuclei, and constitute the sole output of all motor coordination in the cerebellar cortex.<sup>61</sup> The initial

study of Ritvo *et al.* indicated that the number of Purkinje cells in the vermis of the cerebellum was 15 standard deviations below the mean and approximately 8 standard deviations below the mean bilaterally in the cerebral hemispheres of the individuals with autism compared with controls. Vargas *et al.* found that brain tissues of autistic patients showed extensive neuroglial responses characterized by microglial and astroglial activation.<sup>62</sup> In the brains of autistic patients, the most prominent histological changes were observed in the cerebellum, characterized by a patchy loss of neurons in the Purkinje cell layer and granular cell layer in nine out of ten cerebella; one of these cerebella also showed an almost complete loss of Purkinje cells from the Purkinje cell layer, as well as a marked loss of granular cells. Kern and Jones have summarized the important role of Purkinje cell abnormalities in autism, especially the susceptibility of these cells to oxidative stress during GSH depletion.<sup>63</sup> Such depletion can be due to reduction of GSH due to excessive NAPQI exposure and/or to GSH depletion associated with elevated dopamine secondary to dopamine β-hydroxylase inhibition by phenolic *Clostridia* metabolites. The metabolite of acetaminophen, 4-amino phenol, also caused depletion of GSH.<sup>64</sup>

## IMMUNE ABNORMALITIES ASSOCIATED WITH ACETAMINOPHEN USE

An important but under-appreciated aspect of acetaminophen toxicity is that direct, drug-induced harm accounts for only part of the overall syndrome of acetaminophen-induced liver injury. The reason for this is that the initial wave of drug-induced hepatocellular destruction is followed by a robust innate immune response, in which invading inflammatory cells release toxic oxidants and cause a second wave of destruction. The collateral damage inflicted by inflammatory cells can be so severe as to double the degree of tissue injury caused by acetaminophen alone.<sup>34</sup>

Prymula and colleagues compared post-vaccine fever and post-vaccine changes in vaccine-specific

antibody titers in children who either did or did not receive scheduled acetaminophen with immunizations.<sup>37</sup> Four hundred and fifty nine healthy infants undergoing initial and booster vaccinations for a variety of vaccines were randomized to receive or not to receive prophylactic acetaminophen at the time of and for 24 h after vaccinations. Fever was significantly less common in acetaminophen-treated children after initial and booster vaccination. Children who received acetaminophen had significantly lower antibody titers to each vaccine-related antigen, although it was judged that the titers would probably be protective against the diseases for which they had been immunized. In children with severe sulfation deficiency with a predisposition to autism, vaccination without an adequate immune response might lead to viral infection even with the attenuated strains of the viral vaccines due to a lack of GSH needed for effective immune response.<sup>46</sup>

Acetaminophen sales were high in English-speaking countries, and were positively associated with asthma symptoms, eczema and allergic rhinoconjunctivitis in children aged 13–14, and with wheeze, diagnosed asthma, rhinitis and bronchial responsiveness in adults. The prevalence of wheeze increased by 0.52% in 13–14 year olds and by 0.26% in adults ( $p < 0.0005$ ) for each gram increase in per capita acetaminophen sales. Between 1980 and 2003, the prevalence of pediatric asthma in the United States increased from 3.6% to 5.8%, and similar increases were observed throughout the world.<sup>45</sup> It has been speculated that frequent use of acetaminophen might influence asthma and rhinitis by depleting levels of reduced GSH in the nose and airways, thus shifting the oxidant/antioxidant balance in favor of oxidative stress and increasing inflammation.<sup>44</sup>

Acetaminophen decreases GSH levels, principally in the liver and kidneys, but also in the lungs.<sup>46</sup> These decreases are dose dependent; overdose levels of acetaminophen are cytotoxic to pneumocytes and cause acute lung injury, whereas nontoxic, therapeutic doses produce smaller, but significant reductions in GSH levels in type II pneumocytes and alveolar macrophages.<sup>46</sup> Among healthy young volunteers, significantly lower serum antioxidant capacity has been seen within 2 weeks of ingestion of 1 g of acetaminophen.<sup>46</sup> By depleting GSH levels, acetaminophen weakens the ability of

the host to mitigate oxidative stress produced by reactive oxygen species (ROS) such as superoxide anions ( $O_2^-$ ), hydroxyl ( $\bullet OH$ ), and peroxy ( $ROO^-$ ) radicals.<sup>46</sup> Finally, when GSH levels are low, defective processing of disulfide bonds that are key in antigen presentation has been hypothesized.<sup>46</sup> It is conceivable that decreased levels of GSH guide the expression of T-helper cell pathways by altering antigen presentation and recognition, thereby favoring the T2 allergic-dominant pathway. In a study of children with autism spectrum disorders (ASDs) in Sweden, airway symptoms of wheezing and physician-diagnosed asthma in the baseline investigation in infants and toddlers were associated with ASD 5 years later.<sup>52</sup>

Skewed T1 or T2 responses were also indicated in ASD children.<sup>65, 66</sup> Analysis by Jyonouchi *et al.* of adaptive immune responses revealed markedly variable T1/T2 cytokine levels in ASD children compared with control siblings.<sup>67</sup> It has been reported that immune responses in autistic children are relatively skewed to T2 on the basis of intracellular staining of T1/T2 cytokines.<sup>65</sup> Higher levels of IgG, IgA, and IgE allergies to various foods, but especially milk and wheat, have been reported in children with autism, together with abnormal cellular immune responses to milk, wheat, and soy.<sup>67</sup> It has been shown that depleting GSH in brain microglia and astroglia induces a neuroinflammatory response that results in both significant cytokine release and the release of material that is toxic to neurons.

In addition, the ability of GSH to downregulate nuclear factor- $\kappa B$ , and the inverse association between alveolar GSH levels and bronchial responsiveness, suggests that GSH may modify asthma inflammation. Secondly, studies in animals have found that acetaminophen can deplete the lung of GSH. These effects in macrophages raise the possibility that acetaminophen might also influence atopic diseases more generally through another mechanism, namely the promotion of atopy, since depletion of GSH in antigen-presenting cells promotes T-helper cell 2-type cytokine responses. This might explain why, in children, acetaminophen sales were associated with atopic eczema as well as with asthma and rhinitis. As of 2012, 416 studies had demonstrated an association between autism and immune abnormalities and/or inflammation.<sup>3</sup>

The beginning of the rapid increase in autism in around 1980 coincides with the rapid increase in asthma, both of which coincide with the rapid increase in the use of acetaminophen following the Reye's syndrome scare over a possible association with aspirin.

It would seem likely that perhaps some children on the autistic spectrum might have both brain and lung abnormalities caused by acetaminophen. Is there any evidence for such combined abnormalities? Indeed, at the annual meeting of the American College of Chest Physicians in Honolulu, Hawaii in 2011, Barbara Stewart, a pediatric pulmonologist, found that a significant lung abnormality was present in 100% of children ( $n = 47$ ) on the autistic spectrum who were examined, but in none of  $>300$  children without autism.<sup>68</sup> Most of the children with autism had been referred to her clinic due to persistent cough unresponsive to treatment. She noticed during bronchoscopy examinations, in which a lighted tube is inserted into the lungs, that, although the airways of the children initially appeared normal, the lower airway had doubled branches, or "doublets". Dr. Stewart said "when airways divide beyond the first generation, they typically branch like a tree, with one branch on one side and one on the other. A doublet occurs when there are twin branches that come off together instead of one, which are exactly symmetrical, in each of the lower locations that can be seen." In a study of ASD children in Sweden, airway symptoms of wheezing and physician-diagnosed asthma in the baseline investigation in infants and toddlers were associated with ASD 5 years later.<sup>52</sup> It would seem very useful to examine the use of acetaminophen both prenatally and postnatally in the children with the abnormal lung anatomy.

## DNA DAMAGE

As of 2012, 145 studies had shown an association between mitochondria function and autism.<sup>3</sup> The data of Cover *et al.* showed that nitration of mitochondrial proteins and depletion of mitochondrial DNA after acetaminophen overdose in mice was due to peroxynitrite formation.<sup>32</sup> In contrast, nDNA damage was not directly caused by peroxynitrite. Nuclear DNA damage after acetaminophen overdose

is likely to be caused by DNase(s) unrelated to the caspase-activated DNase, which is typically responsible for DNA fragmentation during apoptosis. The data from Cover *et al.* suggest that the activation of these DNase(s) is dependent on the mitochondrial oxidant stress and peroxynitrite formation.<sup>32</sup>

Recent data have demonstrated that nitrated tyrosine residues, as well as acetaminophen adducts, occur in the necrotic cells following toxic doses of acetaminophen. Nitrotyrosine was postulated to be mediated by peroxynitrite, a reactive nitrogen species formed by the very rapid reaction of superoxide and nitric oxide (NO). Peroxynitrite is normally detoxified by GSH, which is depleted in acetaminophen toxicity. NO synthesis (serum nitrate plus nitrite) was dramatically increased following acetaminophen.

## NAPQI

The metabolism of acetaminophen (Figure 2) by CYP2E1 forms NAPQI, a reactive metabolite that binds to cysteine residues in cellular proteins and forms acetaminophen protein adducts, referred to as acetaminophen-cysteine complexes (APAP-CYS).

Heard *et al.* demonstrated that low concentrations of APAP-CYS are detectable in serum following therapeutic dosing with acetaminophen in the vast majority of individuals.<sup>49</sup> The APAP-CYS concentrations following acetaminophen overdose varied widely. Serum APAP-CYS concentrations in patients with hepatic injury following acetaminophen overdose were generally much higher than those observed during therapeutic dosing. However, three overdose patients had APAP-CYS concentrations that were of the same order of magnitude as those observed with therapeutic dosing. Adduct concentrations varied according to the degree of exposure. Currently, an absolute adduct level exceeding  $1.1 \mu\text{mol/L}$  appears consistent with acetaminophen toxicity. While therapeutic dosing generally produces concentrations below  $0.5 \mu\text{mol/L}$ , values in people taking recommended dosages have been reported to be very close to the lower limit of APAP-CYS concentration associated with acetaminophen toxic overdoses ( $1.0 \mu\text{mol/L}$ ).

## NAPQI AND ANOMALOUS HAIR METAL VALUES IN AUTISM

Concentrations of the NAPQI metabolite increased to values as high as 1.0  $\mu\text{mol/L}$  or 1000 nmol/L serum following therapeutic doses of acetaminophen. Hair proteins contain a considerable amount of cysteine, and hair proteins in hair follicles and would be expected to react with NAPQI to form NAPQI adducts with cysteine sulfhydryl groups in hair. Such cysteine groups also react strongly with heavy metals such as mercury. In comparison, the mean blood mercury concentration of children with autism was 19.53 nmol/L. Thus the concentration of NAPQI might be as much as 51 times the concentration of total mercury. Since NAPQI amounts are so much higher than mercury values, it would be expected that NAPQI might significantly reduce the capacity of hair proteins to bind mercury if the binding capacity of the hair sulfhydryl groups was exceeded by the previous binding of NAPQI. Such inhibition by NAPQI may help to explain anomalous data in the measurement of mercury in hair samples of children with autism. In two baby hair studies in which samples were obtained at first haircuts, mercury values in hair were much lower in children with autism compared with those of normal controls.<sup>69, 70</sup> In addition, children with the most severe symptoms of autism had the lowest amount of hair mercury. Studies in Kuwait and Saudi Arabia found opposite results in children with autism; children with autism had much higher amounts of hair mercury as well as other heavy metals, but in both of these studies the children were much older (mean ages 8.8 years in Saudi Arabia and 4.2 years in Kuwait) than those in the baby hair studies (12–24 months of age).<sup>71</sup> <sup>72</sup> All of this data could be explained by patterns of acetaminophen drug use in children. Acetaminophen is commonly used prophylactically to prevent fever in infants and toddlers who receive the bulk of their vaccines in the first 2 years of life. In contrast, few vaccines and attendant prophylactic acetaminophen are administered to older children with autism in the age range in which hair mercury values were elevated compared with controls. In these older children, hair metals such as mercury would be considerably higher due to a lack of competition from NAPQI for reactive sulfhydryl groups in hair proteins in the hair follicles.

## SPECIAL CONCERNS ABOUT THE LONG-TERM DEFECTIVE QUALITY CONTROL OF ACETAMINOPHEN PRODUCTS

All of the information to this point has been directed to the toxicity of pure acetaminophen. Toxicity is usually due to exceeding the recommended daily dosage, often occurring due to accidental combination use of multiple products containing acetaminophen or through changes in dosage amounts by manufacturers. Adding to this are reports that acetaminophen products have been contaminated at various times, potentially increasing health risks. These problems have been documented in Food and Drug Administration inspection reports and subsequent recalls.

From the time it was introduced in 1955, acetaminophen has become one of the most successful OTC drugs, and has earned the primary manufacturer, Johnson & Johnson, an estimated US\$1.3 billion every year.<sup>73</sup> Within the last 10 years, the FDA has chronicled manufacturing problems including mislabeling of children's tablets, where packages of the product listed an incorrect amount of the ingredients per tablet, inadequate cleaning of equipment used for manufacturing, insufficient follow-up in investigating consumer complaints, product mix-ups, and contamination with metal fragments and bacteria.<sup>74</sup> <sup>75</sup> The latter resulted in a recall of 135 million bottles of children's medications containing acetaminophen.

Additional recalls occurred in 2009 and 2010 due to trace amounts of the chemical 2,4,6-tribromoanisole found in infant, children, and adult OTC products. This recall was prompted by 775 consumer reports of nausea, vomiting, stomach pains and diarrhea received by FDA due to the contamination.<sup>76</sup> It was found that the chemical 2,4,6-tribromoanisole (a metabolite of a chemical fungicide tribromophenol) was being used in a manufacturing facility to treat the wood used in pallets used for transporting packaged materials.<sup>77</sup> Tribromoanisole is produced when naturally-occurring airborne fungi and/or bacteria (usually *Aspergillus* sp., *Penicillium* sp., *Actinomyces*, *Botrytis cinerea*, *Rhizobium* sp., or *Streptomyces*) are presented with brominated



phenolic compounds, which they then convert into bromoanisole derivatives.

Other FDA-publicized inspection reports of acetaminophen-manufacturing facilities have listed inadequate quality controls, lack of safeguards to isolate “rejected” raw materials and other drugs, and several human errors resulting in product mix-ups.<sup>78</sup> Several recalls have occurred due to failed FDA manufacturing inspections and contaminated products affecting >300 million bottles of adult and children’s medicines.<sup>79–82</sup> Manufacturing controls are an integral aspect of pharmaceutical production to ensure public safety, and disregard or deficiency can lead to serious outcomes. In response to the high number of inadequacies found through inspections and numerous recalls, US health authorities have taken over supervision of three acetaminophen-manufacturing plants to mitigate risks.<sup>83</sup>

## A CALL TO ACTION

A large-scale, long-term study of both prenatal and postnatal effects of acetaminophen exposure on the incidence of autism, attention deficit with hyperactivity, and asthma should be conducted. However, such a study would probably take at least 5 years. If acetaminophen is indeed the cause of all of these illnesses, should acetaminophen use continue for such a long period of time while millions of additional children are further affected? Respected physicians consider that the connection of acetaminophen with asthma has been proven beyond a reasonable doubt. Dr McBride, Professor of Pediatrics at Department of Pediatrics, Northeast Ohio Medical University, Rootstown, Ohio summarizes the evidence for the acetaminophen asthma findings<sup>45</sup>: “There remains a possibility that confounding variables might explain some or all of the association between acetaminophen and asthma. For this reason we need further studies. At present, however, I need further studies not to prove that acetaminophen is dangerous but, rather, to prove that it is safe. Until such evidence is forthcoming, I will recommend avoidance of acetaminophen by all children with asthma or those at risk for asthma and will work to make patients, parents, and primary care

providers aware of the possibility that acetaminophen is detrimental to children with asthma.”

Since *all children* may be at risk from asthma, Dr McBride is in effect saying that acetaminophen is contraindicated for the treatment of *any* children. Although the case for acetaminophen being a cause of autism and attention deficit with hyperactivity may not be as strong as the case for asthma, the severe asthma risk combined with the risks of autism and attention deficit with hyperactivity are so severe that we as a society should maintain a degree of caution with acetaminophen given the proven overall toxicity due to accidental overdose of the drug, and the availability of ibuprofen or abstaining from treatment as alternatives. A large-scale trial of acetaminophen restriction in pregnancy and the first 3 years of life is warranted to test the hypothesis that acetaminophen is a causative agent in autism, asthma, and attention deficit with hyperactivity. Due to the increased risks associated with accidental overuse of acetaminophen, increased public awareness of such risks is paramount.

Given associations of acetaminophen with increased rates of cancer, increased testicular damage, increased rates of asthma, and allergy, plausible causation of autism, and *in vitro* evidence of brain damage associated with metabolites of acetaminophen, new assessments of the relative risk of aspirin causing Reye’s syndrome versus the risks of acetaminophen in children should be undertaken. If a clear link between acetaminophen use pre- or post-natally and autism is established, medical practice guidelines may need to be adjusted and alternative analgesics or antipyretics, such as ibuprofen, recommended. However, such a relationship may be difficult to establish, as studies may have to account for variations in dosage amounts, formats, combination product use, and for the possibility of product recalls given the recent reported manufacturing issues surrounding acetaminophen.

## NOTE ADDED IN PROOF

After this article was submitted for publication on Lcp04."4235, the following article was published prior to publication which strongly supports the hypothesis in the current article.



In 2013, Bauer and Kriebel reported that prenatal use of acetaminophen was strongly correlated with autism/Autism Spectrum Disorder prevalence ( $r = 0.80$ ) using all available country-level data ( $n = 8$ ) for the period 1984 to 2005. In addition, the authors found that after acetaminophen became

commonly used to treat circumcision pain after 1995, there was a strong correlation between country-level ( $n = 9$ ) autism/ASD prevalence in males and a country's circumcision rate ( $r = 0.98$ ). A very similar pattern was seen among US states and when comparing the three main racial/ethnic groups in the US.<sup>84</sup>

## REFERENCES

- Kim YS, Leventhal BL, Koh YJ, Fombonne E, Laska E, Lim EC, *et al.* Prevalence of autism spectrum disorders in a total population sample. *Am J Psychiatry*. 2011;168(9):904–12.
- Hertz-Picciotto I, Delwiche L. The rise in autism and the role of age at diagnosis. *Epidemiology*. 2009;20(1):84–90.
- Rossignol DA, Frye RE. A review of research trends in physiological abnormalities in autism spectrum disorders: Immune dysregulation, inflammation, oxidative stress, mitochondrial dysfunction and environmental toxicant exposures. *Mol Psychiatry*. 2012;17(4):389–401.
- Shaw W. The unique vulnerability of the human brain to toxic chemical exposure and the importance of toxic chemical evaluation and treatment in orthomolecular psychiatry. *J Orthomol Med*. 2010;25(3):125–34.
- Miller MT. Thalidomide embryopathy: A model for the study of congenital incontinent horizontal strabismus. *Trans Am Ophthalmol Soc*. 1991;89:623–74.
- Disability-analysis of State Reports, reviewed by the Committee on the Rights of the Child (CRC) in its Pre-Sessional Working Group, 55<sup>th</sup> Session. International Disability Alliance 2010;8.
- <http://www.autism-society.org/about-autism/>. Autism Society Website 2012. Accessed August 31, 2013.
- Reed G, Galindo MA. Cuba's National Immunization Program. *MEDICC Review*. 2008;9(1):5–7.
- Bäckström M, Hägg S, Mjörndal T, Dahlqvist R. Utilization pattern of metamizole in northern Sweden and risk estimates of agranulocytosis. *Pharmacoepidemiol Drug Saf*. 2002;11(3):239–45.
- Torres AR. Is fever suppression involved in the etiology of autism and neurodevelopmental disorders? *BMC Pediatr*. 2003;2:3–9.
- Schultz ST, Klonoff-Cohen HS, Wingard DL, Akshoomoff NA, Macera CA, Ji M. Acetaminophen (paracetamol) use, measles-mumps rubella vaccination, and autistic disorder: The results of a parent survey. *Autism*. 2008;12(3):293–307.
- Schultz ST. Can autism be triggered by acetaminophen activation of the endocannabinoid system? *Acta Neurobiol Exp*. 2010;70(2):227–31.
- Schultz S, Desilva M, Gu TT, Qiang M, Whang K. Effects of the analgesic acetaminophen (paracetamol) and its para-aminophenol metabolite on viability of mouse-cultured cortical neurons. *Basic Clin Pharmacol Toxicol*. 2011;110(2):141–4.
- Becker KG, Schultz ST. Similarities in features of autism and asthma and a possible link to acetaminophen use. *Med Hypotheses*. 2010;74(1):7–11.
- Posadas I, Santos P, Blanco A, Muñoz-Fernández M, Ceña V. Acetaminophen induces apoptosis in rat cortical neurons. *PLoS ONE*. 2010;15(12):e15360.
- U.S. Food and Drug Administration. For consumers – Don't double up on acetaminophen. 2013. Available at: <http://www.fda.gov/ForConsumers/ConsumerUpdates/ucm336581.htm>. Accessed March 19, 2013.
- Connor DF. Problems of overdiagnosis and overprescribing in ADHD. *Psychiatric Times*. 2011;28(8):1–8.
- Center For Disease Control. Mental health in the United States: Prevalence of diagnosis and medication treatment for attention-deficit/hyperactivity disorder – United States, 2003. *MMWR*. 2005;54(34):842–7.
- Safer DJ. Are stimulants overprescribed for youths with ADHD? *Ann Clin Psychiatry*. 2000;12(1):55–62.
- Safer D, Krager JM. Hyperactivity and inattentiveness. School assessment of stimulant treatment. *Clin Pediatr*. 1989;28(5):216–21.
- Safer DJ, Zito JM, Fine EM. Increased methylphenidate usage for attention deficit disorder in the 1990s. *Pediatrics*. 1996;98(6):1084–8.
- Safer DJ, Krager JM. The increased rate of stimulant treatment for hyperactive/inattentive students in secondary schools. *Pediatrics*. 1994;94(4):462–4.
- Safer DJ, Malever M. Stimulant treatment in Maryland Public Schools. *Pediatrics*. 2009;106(3):533–9.
- Safer DJ, Krager JM. A survey of medication treatment for hyperactive/inattentive students. *J Am Med Assoc*. 1988;260(15):2256–8.
- Mandell DS, Thompson WW, Weintraub ES, Destefano F, Blank MB. Trends in diagnosis rates for autism and ADHD at hospital discharge in the context of other psychiatric diagnoses. *Psychiatr Serv*. 2005;56(1):56–62.

26. Orłowski JP, Hanhan UA, Fiallos MR. Is aspirin a cause of Reye's syndrome? A case against. *Drug Saf.* 2002;25(4):225–31.
27. Van Bever HP, Quek SC, Lim T. Aspirin, Reye syndrome, Kawasaki disease, and allergies; a reconsideration of the links. *Br Med J.* 2004;89(12):1178.
28. Weiner DL. Reye Syndrome. 2011. Available at: <http://EMedicine.Medscape.com>. Accessed August 31, 2013.
29. Shaw W. Possible role of lysolecithins and nonesterified fatty acids in the pathogenesis of Reye's syndrome, sudden infant death syndrome, acute pancreatitis, and diabetic ketoacidosis. *Clin Chem.* 1985;31(7):1109–15.
30. Kristensen DM, Hass U, Lesné L, Lottrup G, Jacobsen PR, Desdoits-Lethimonier C, et al. Intrauterine exposure to mild analgesics is a risk factor for development of male reproductive disorders in human and rat. *Human Reprod.* 2010;26(1):235–44.
31. Adjei AA, Gaedigk A, Simon SD, Weinshilboum RM, Leeder JS. Interindividual variability in acetaminophen sulfation by human fetal liver: Implications for pharmacogenetic investigations of drug-induced birth defects. *Birth Defects Res A Clin Mol Teratol.* 2008;82(3):155–65.
32. Cover C, Mansouri A, Knight TR, Bajt ML, Lemasters JJ, Pessayre D, et al. Peroxynitrite-induced mitochondrial and endonuclease-mediated nuclear DNA damage in acetaminophen hepatotoxicity. *J Pharmacol Exp Ther.* 2005;315(2):879–87.
33. Ruepp SU, Tonge RP, Shaw J, Wallis N, Pognan F. Genomics and proteomics analysis of acetaminophen toxicity in mouse liver. *Toxicol Sci.* 2012;65(1):135–50.
34. Maher JJ. DAMPs ramp up drug toxicity. *J Clin Invest.* 2009;119(2):246–9.
35. Lee SS, Buters JT, Pineau T, Fernandez-Salguero P, Gonzalez FJ. Role of CYP2E1 in the hepatotoxicity of acetaminophen. *J Biol Chem.* 1996;271(20):12063–7.
36. Yamaura K, Ogawa K, Yonekawa T, Nakamura T, Yano S, Ueno K. Inhibition of the antibody production by acetaminophen independent of liver injury in mice. *Biol Pharm Bull.* 2002;25(2):201–5.
37. Prymula R, Siegrist CA, Chlibek R, Zemlickova H, Vackova M, Smetana J, et al. Effect of prophylactic paracetamol administration at time of vaccination on febrile reactions and antibody responses in children: Two open-label, randomised controlled trials. *Lancet.* 2009;374:1339–50.
38. Homme JH, Fischer PR. Prophylactic paracetamol at the time of infant vaccination reduces the risk of fever but also reduces antibody response. *Evid Based Med.* 2010;15(2):50–1.
39. Fennes AZ, Kirkpatrick III HM, Patel VV, Sweetman L, Emmett M. Increased anion gap metabolic acidosis as a result of 5-oxoproline (pyroglutamic acid): A role for acetaminophen. *Clin J Am Soc Nephrol.* 2006;1(3):441–7.
40. Pitt JJ, Brown GK, Clift V, Christodoulou J. Atypical pyroglutamic aciduria: Possible role of paracetamol. *J Inherit Metab Dis.* 1990;13(5):755–6.
41. Miller AL. Liver damage from chronic acetaminophen dosing is dangerous, but not the only risk. *Alt Med Rev.* 2009;14(4):322–3.
42. Franciscus A, Highleyman L. Acetaminophen and your liver. *HCSP Fact Sheet.* 2009;2.2:1–3.
43. Walter RB, Milano F, Brasky TM, White E. Long-term use of acetaminophen, aspirin, and other nonsteroidal anti-inflammatory drugs and risk of hematologic malignancies: results from the prospective vitamins and lifestyle (VITAL) study. *J Clin Oncol.* 2011;94:2424–2431.
44. Newson RB, Shaheen SO, Chinn S, Burney PG. Paracetamol sales and atopic disease in children and adults: An ecological analysis. *Eur Respir J.* 2000;16(5):817–23.
45. McBride JT. Association of acetaminophen and asthma prevalence and severity. *Pediatrics.* 2011;128(6):1181–5.
46. Eneli I, Sadri K, Camargo C Jr, Barr RG. Acetaminophen and the risk of asthma: The epidemiologic and pathophysiologic evidence. *Chest.* 2005;127(2):604–12.
47. Somers GF. Pharmacological properties of Thalidomide (α-phthalimido glutarimide), a new sedative hypnotic drug. *Br J Pharmacol Chemother.* 1960;15:111–116.
48. Good P. Did acetaminophen provoke the autism epidemic? *Alt Med Rev.* 2009;14(4):364–72.
49. Heard KJ, Green JL, James LP, Judge BS, Zolot L, Rhyee S, et al. Acetaminophen-cysteine adducts during therapeutic dosing and following overdose. *BMC Gastroenterol.* 2011;11:20.
50. Krakowiak P, Walker CK, Bremer AA, Baker AS, Ozonoff S, Hansen RL, et al. Maternal metabolic conditions and risk for autism and other neurodevelopmental disorders. *Pediatrics.* 2012;129(5):e1121–8.
51. Windham GC, Zhang L, Gunier R, Croen LA, Grether JK. Autism spectrum disorders in relation to distribution of hazardous air pollutants in the San Francisco Bay area. *Environ Health Perspect.* 2006;114(9):1438–44.
52. Larsson M, Weiss B, Janson S, Sundell J, Bornehag CG. Associations between indoor environmental factors and parental-reported autistic spectrum disorders in children 6–8 years of age. *Neurotoxicology.* 2009;30(5):822–31.
53. Kim SJ, Lee MY, Kwon do Y, Kim SY, Kim YC. Alteration in metabolism and toxicity of acetaminophen upon repeated administration in rats. *J Pharmacological Sci.* 2009;111(2):175–81.
54. Alcorn J, McNamara PJ. Pharmacokinetics in the newborn. *Adv Drug Deliv Rev.* 2003;55(5):667–86.
55. Högestätt ED, Jönsson BA, Ermund A, Andersson DA, Björk H, Alexander JP, et al. Conversion of acetaminophen to the bioactive N-acetylphenolamine AM404 via fatty acid amide hydrolase-dependent arachidonic acid conjugation in the nervous system. *J Biol Chem.* 2005;280(36):31405–12.

56. O'Reilly BA, Waring RH. Enzyme and sulphur oxidation deficiencies in autistic children with known food/chemical intolerances. *J Orthomol Med.* 1993;8(4):198–200.
57. Alberti A, Pirrone P, Elia M, Waring RH, Romano C. Sulfation deficit in “low functioning” autistic children: A pilot study. *Biol Psychiatry.* 1999;46(3):420–4.
58. Schlune A, Mayatepek E. Glutathione synthetase deficiency – An inborn error of the gamma-glutamyl cycle. *J Pediatr Sci.* 2011;3(70):2505–9.
59. Adams JB, Baral M, Geis E, Mitchell J, Ingram J, Hensley A, *et al.* The severity of autism is associated with toxic metal body burden and red blood cell glutathione levels. *J Toxicol.* 2009;2009:532640.
60. Ritvo ER, Freeman BJ, Scheibel AB, Duong T, Robinson H, Guthrie D, *et al.* Lower Purkinje cell counts in the cerebella of four autistic subjects: Initial findings of the UCLA-NSAC autopsy research reports. *Am J Psychiatry.* 1986;143(7):862–6.
61. Purves D, Augustine GJ, Fitzpatrick D, Hall WC, LaMantia AS, McNamara JO, *et al.* *Neuroscience*, 4th edn. Sunderland, MA: Sinaur Associates; 2008:432–4.
62. Vargas DL, Nascimbene C, Krishnan C, Zimmerman AW, Pardo CA. Neuroglial activation and neuroinflammation in the brain of patients with autism. *Ann Neurol.* 2005;57(1):67–81.
63. Kern JK, Jones AM. Evidence of toxicity, oxidative stress, and neuronal insult in autism. *J Toxicol. Environ Health B Crit Rev.* 2006;9(6):485–99.
64. Fu X, Chen TS, Ray MB, Nagasawa HT, Williams WM. p-Aminophenol-induced hepatotoxicity in hamsters: Role of glutathione. *J Biochem Mol Toxicol.* 2004;18(3):154–61.
65. Gupta S, Aggarwal S, Roshanravan B, Lee T. Th1- and Th2-like cytokines in CD4+ and CD8+ T cells in autism. *J Neuroimmunol.* 1998;85(1):106–9.
66. Singh VK. Plasma increase of interleukin-12 and interferon-gamma. Pathological significance in autism. *J Neuroimmunol.* 1996;66(1-2):143–5.
67. Jyonouchi H, Sun S, Le H. Proinflammatory and regulatory cytokine production associated with innate and adaptive immune responses in children with autism spectrum disorders and developmental regression. *J Neuroimmunol.* 2001;120(1-2):170–9.
68. Harrison L. Autism linked to unusual shapes in lungs study shows children born with a unique airway shape may have autism. *WebMD Health News.* 2011. Available at: [http://www.webmd.com/brain/autism/news/20111025/autism-linked-to-unusual-airway-shapes?src=RSS\\_PUBLC](http://www.webmd.com/brain/autism/news/20111025/autism-linked-to-unusual-airway-shapes?src=RSS_PUBLC). Accessed August 31, 2013.
69. Holmes AS, Blaxill MF, Haley BE. Reduced levels of mercury in first baby haircuts of autistic children. *Int J Toxicol.* 2003;22(4):277–85.
70. Adams JB, Romdalvik J, Levine KE, Hu LW. Mercury in first-cut baby hair of children with autism versus typically-developing children. *Toxicol Environ Chem.* 2008;90(4):739–53.
71. Fido A, Al-Saad S. Toxic trace elements in the hair of children with autism. *Autism.* 2005;9(3):290–8.
72. Al-Ayadhi LY. Heavy metals and trace elements in hair samples of autistic children in central Saudi Arabia. *Neurosciences.* 2005;10(3):213–8.
73. Easton T, Herrera S. Despite bad publicity and costly legal settlements, Johnson & Johnson refuses to put ample warnings on its Tylenol labels. Torah View 2012. Available at: <http://www.torahview.com/bris/html/tylenol.html>. Accessed April 26, 2012.
74. Kavilanz P. Tylenol plant: From bad to worse. CNN Money 2010. Available at: [http://money.cnn.com/2010/06/22/news/companies/tylenol\\_plant\\_inspection\\_history/index.htm](http://money.cnn.com/2010/06/22/news/companies/tylenol_plant_inspection_history/index.htm). Accessed April 26, 2012.
75. Kavilanz P. Tylenol plant still plagued by FDA violations. CNN Money 2012. Available at: [http://money.cnn.com/2010/12/01/news/companies/tylenol\\_plant\\_new\\_problems/index.htm](http://money.cnn.com/2010/12/01/news/companies/tylenol_plant_new_problems/index.htm). Accessed April 26, 2012.
76. Kavilanz P. Tylenol recall: Serious side effects investigated. CNN Money 2010. Available at: [http://money.cnn.com/2010/05/25/news/companies/tylenol\\_recall\\_adverse\\_consumer\\_complaints/index.htm](http://money.cnn.com/2010/05/25/news/companies/tylenol_recall_adverse_consumer_complaints/index.htm). Accessed April 26, 2012.
77. Kavilanz P. New recalled Tylenol linked to shut plant. CNN Money 2010. Available at: [http://money.cnn.com/2010/10/19/news/companies/new\\_tylenol\\_recall/index.htm](http://money.cnn.com/2010/10/19/news/companies/new_tylenol_recall/index.htm). Accessed April 26, 2012.
79. Baker SL. Drugs like Tylenol can be contaminated with mold and chemicals. NaturalNews Com 2010. Available at: [http://www.naturalnews.com/027965\\_Tylenol\\_recall.html](http://www.naturalnews.com/027965_Tylenol_recall.html). Accessed April 26, 2012.
80. Heavey S. FDA finds grime at J&J plant, urges use of generics. Reuters 2010. Available at: <http://www.reuters.com/article/2010/05/05/us-johnsonandjohnson-recall-idUSTRE64367Z20100505>. Accessed April 26, 2012.
81. Silverman E. Which Drugmaker Fails Most FDA Inspections? Pharmalot com 2011. Available at: <http://www.pharmalot.com/2011/03/which-drugmaker-fails-most-fda-inspections/>. Accessed April 26, 2012.
82. Krauskopf L. More Tylenol Recalled. The Huffington Post 2011. Available at: [http://www.huffingtonpost.com/2011/03/30/tylenol-recalled-\\_n\\_842450.html](http://www.huffingtonpost.com/2011/03/30/tylenol-recalled-_n_842450.html). Accessed April 27, 2012.
83. Kavilanz P. U.S. takes over three Tylenol plants. CNN Money 2011. Available at: [http://money.cnn.com/2011/03/10/news/companies/johnson\\_mccneil\\_fda\\_action/index.htm](http://money.cnn.com/2011/03/10/news/companies/johnson_mccneil_fda_action/index.htm). Accessed April 27, 2012.
84. Bauer AZ, Kriebel D. Prenatal and perinatal analgesic exposure and autism: an ecological link. *Environ Health.* 2013;12:41. doi: 10.1186/1476-069X-12-41.